

Now let us do some...

# Counting

(Chapter 6)

Oct 1, 2015

CS 320

1

## Basic Counting Principles

**Counting problems** are of the following kind:

“**How many** different 8–letter passwords are there?”

“**How many** possible ways are there to pick 11 soccer players out of a 20–player team?”

Most importantly, counting is the basis for computing **probabilities of discrete events**.

(“What is the probability of winning the lottery?”)

Oct 1, 2015

CS 320

2

## Basic Counting Principles

### The sum rule:

If a task can be done in  $n_1$  ways and a second task in  $n_2$  ways, and if these two tasks cannot be done at the same time, then there are  $n_1 + n_2$  ways to do either task.

### Example:

The department will award a free computer to either a CS student or a CS professor. How many different choices are there, if there are 530 students and 15 professors?

There are  $530 + 15 = 545$  choices.

Oct 1, 2015

CS 320

3

## Basic Counting Principles

### Generalized sum rule:

If we have tasks  $T_1, T_2, \dots, T_m$  that can be done in  $n_1, n_2, \dots, n_m$  ways, respectively, and no two of these tasks can be done at the same time, then there are  $n_1 + n_2 + \dots + n_m$  ways to do one of these tasks.

Oct 1, 2015

CS 320

4

## Basic Counting Principles

### The product rule:

Suppose that a procedure can be broken down into two successive tasks. If there are  $n_1$  ways to do the first task and  $n_2$  ways to do the second task after the first task has been done, then there are  $n_1 n_2$  ways to do the procedure.

### Generalized product rule:

If we have a procedure consisting of sequential tasks  $T_1, T_2, \dots, T_m$  that can be done in  $n_1, n_2, \dots, n_m$  ways, respectively, then there are  $n_1 \cdot n_2 \cdot \dots \cdot n_m$  ways to carry out the procedure.

Oct 1, 2015

CS 320

5

## Basic Counting Principles

### Example:

How many different license plates are there that contain exactly three English letters ?

### Solution:

There are 26 possibilities to pick the first letter, then 26 possibilities for the second one, and 26 for the last one.

So there are  $26 \cdot 26 \cdot 26 = 17576$  different license plates.

Oct 1, 2015

CS 320

6

## Basic Counting Principles

The sum and product rules can also be phrased in terms of **set theory**.

**Sum rule:** Let  $A_1, A_2, \dots, A_m$  be disjoint sets. Then the number of ways to choose any element from one of these sets is  $|A_1 \cup A_2 \cup \dots \cup A_m| = |A_1| + |A_2| + \dots + |A_m|$ .

**Product rule:** Let  $A_1, A_2, \dots, A_m$  be finite sets. Then the number of ways to choose one element from each set independently in the order  $A_1, A_2, \dots, A_m$  is  $|A_1 \times A_2 \times \dots \times A_m| = |A_1| \cdot |A_2| \cdot \dots \cdot |A_m|$ .

Oct 1, 2015

CS 320

7

## Inclusion-Exclusion

How many bit strings of length 8 start with a 1 or end with 00?

**Task 1:** Create a string of length 8 that starts with a 1.

There is one way to pick the first bit (1).

For each of these there are two ways to pick the second bit (0 or 1).

For each of these, two ways to pick the third bit (0 or 1),

⋮

For each, two ways to pick the eighth bit (0 or 1).

**Product rule:** Task 1 can be done in  $1 \cdot 2^7 = 128$  ways.

Oct 1, 2015

CS 320

8

## Inclusion-Exclusion

**Task 2:** Construct a string of length 8 that ends with 00.

There are two ways to pick the first bit (0 or 1), two ways to pick the second bit (0 or 1),

⋮

⋮

two ways to pick the sixth bit (0 or 1),

one way to pick the seventh bit (0), and

one way to pick the eighth bit (0).

**Product rule:** Task 2 can be done in  $2^6 = 64$  ways.

Oct 1, 2015

CS 320

9

## Inclusion-Exclusion

Since there are 128 ways to do Task 1 and 64 ways to do Task 2, does this mean that there are 192 bit strings either starting with 1 or ending with 00?

No, because here Task 1 and Task 2 can be done **at the same time**.

When we carry out Task 1 and create strings starting with 1, some of these strings end with 00.

Therefore, we sometimes do Tasks 1 and 2 at the same time, so **the sum rule does not apply**.

Oct 1, 2015

CS 320

10

## Inclusion-Exclusion

If we want to use the sum rule in such a case, we have to subtract the cases when Tasks 1 and 2 are done at the same time.

How many cases are there, that is, how many strings start with 1 **and** end with 00?

There is one way to pick the first bit (1), two ways for the second, ..., sixth bit (0 or 1), one way for the seventh, eighth bit (0).

**Product rule:** In  $2^5 = 32$  cases, Tasks 1 and 2 are carried out at the same time.

Oct 1, 2015

CS 320

11

## Inclusion-Exclusion

Since there are 128 ways to complete Task 1 and 64 ways to complete Task 2, and in 32 of these cases Tasks 1 and 2 are completed at the same time, there are

$$128 + 64 - 32 = 160 \text{ ways to do either task.}$$

In set theory, this corresponds to sets  $A_1$  and  $A_2$  that are **not** disjoint. Then we have:

$$|A_1 \cup A_2| = |A_1| + |A_2| - |A_1 \cap A_2|$$

This is called the **principle of inclusion-exclusion**.

Oct 1, 2015

CS 320

12

## Inclusion-Exclusion

When counting the number of points in a finite union of sets, we add the points in all the sets. If intersections are non empty we have overcounted, and subtract the number of points in the intersections to compensate. If there are more than three sets we have usually overcompensated and have to add back the number of points in the intersections of three of the sets. This process continues...

See **Theorem 1, page 556**. The notation is hard to put in powerpoint.

Oct 1, 2015

CS 320

13

## Inclusion-Exclusion

We'll draw a Venn diagram for the inclusion-exclusion theorem for  $n = 3$  and  $n = 4$ .

The general case is trickier to prove but you can use the proof on p. 556 once you understand combinations and have done the Binomial Theorem.

Oct 1, 2015


CS 320

14

## Tree Diagrams

How many bit strings of length four do not have two consecutive 1s?

Task 1 Task 2 Task 3 Task 4  
(1<sup>st</sup> bit) (2<sup>nd</sup> bit) (3<sup>rd</sup> bit) (4<sup>th</sup> bit)



There are 8 strings.

Oct 1, 2015

CS 320

15

## The Pigeonhole Principle

**The pigeonhole principle:** If  $(k + 1)$  or more objects are placed into  $k$  boxes, then there is at least one box containing two or more of the objects.

**Example 1:** If there are 11 players in a soccer team that wins 12-0, there must be at least one player in the team who scored at least twice.

**Example 2:** If you have 6 classes from Monday to Friday, there must be at least one day on which you have at least two classes.

Oct 1, 2015

CS 320

16

## The Pigeonhole Principle

**The generalized pigeonhole principle:** If  $N$  objects are placed into  $k$  boxes, then there is at least one box containing at least  $\lceil N/k \rceil$  of the objects.

**Proof:** If every box holds fewer than  $N/k$  objects then there are fewer than  $k(N/k) = N$  objects overall. Thus some box must hold at least  $N/k$  objects, thus  $\lceil N/k \rceil$ , since each box holds an integral number of objects.

**Example 1:** In a 60-student class, at least 12 students will get the same letter grade (A, B, C, D, or F).

**Example 2:** In a 61-student class, at least 13 students will get the same letter grade.

Oct 1, 2015

CS 320

17

## The Pigeonhole Principle

**Example 3:** Assume you have a drawer containing a random distribution of a dozen brown socks and a dozen black socks. It is dark, so how many socks do you have to pick to be sure that among them there is a matching pair?

There are two types of socks, so if you pick at least 3 socks, there must be either at least two brown socks or at least two black socks.

Generalized pigeonhole principle:  $\lceil 3/2 \rceil = 2$ .

Oct 1, 2015

CS 320

18

## Permutations and Combinations

How many different sets of 3 people can we pick from a group of 6?

There are 6 choices for the first person, 5 for the second one, and 4 for the third one, so are there  $6 \cdot 5 \cdot 4 = 120$  ways to do this?

This is not the correct result!

For example, picking person C, then person A, and then person E leads to the **same group** as first picking E, then C, and then A.

However, these cases are counted **separately** in the above equation.

Oct 1, 2015

CS 320

19

## Permutations and Combinations

So how can we compute how many different subsets of people can be picked (that is, we want to disregard the order of picking) ?

To find out about this, we need to look at **permutations**.

A **permutation** of a set of distinct objects is an ordered arrangement of these objects.

An ordered arrangement of  $r$  elements of a set is called an  **$r$ -permutation**.

Oct 1, 2015

CS 320

20

## Permutations and Combinations

**Example:** Let  $S = \{1, 2, 3\}$ .

The arrangement 3, 1, 2 is a permutation of  $S$ .  
The arrangement 3, 2 is a 2-permutation of  $S$ .

The number of  $r$ -permutations of a set with  $n$  distinct elements is denoted by  **$P(n, r)$** .

We can calculate  $P(n, r)$  with the product rule:  
 $P(n, r) = n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot (n-r+1)$ .  
( $n$  choices for the first element,  $(n-1)$  for the second one,  $(n-2)$  for the third one...)

Oct 1, 2015

CS 320

21

## Permutations and Combinations

**Example:**

$$P(8, 3) = 8 \cdot 7 \cdot 6 = 336 \\ = (8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1) / (5 \cdot 4 \cdot 3 \cdot 2 \cdot 1)$$

**General formula:**

$$P(n, r) = n! / (n-r)!$$

Knowing this, we can return to our initial question:  
How many different sets of 3 people can we pick from a group of 6?

Oct 1, 2015

CS 320

22

## Permutations and Combinations

An  **$r$ -combination** of elements of a set is an unordered selection of  $r$  elements from the set. Thus, an  $r$ -combination is simply a subset of the set with  $r$  elements.

**Example:** Let  $S = \{1, 2, 3, 4\}$ .  
Then  $\{1, 3, 4\}$  is a 3-combination from  $S$ .

The number of  $r$ -combinations of a set with  $n$  distinct elements is denoted by  **$C(n, r)$** .

**Example:**  $C(4, 2) = 6$ , since, for example, the 2-combinations of  $\{1, 2, 3, 4\}$  are  $\{1, 2\}$ ,  $\{1, 3\}$ ,  $\{1, 4\}$ ,  $\{2, 3\}$ ,  $\{2, 4\}$ ,  $\{3, 4\}$ .

Oct 1, 2015

CS 320

23

## Permutations and Combinations

How can we calculate  $C(n, r)$ ?

Consider that we can obtain the  $r$ -permutation of a set in the following way:

**First**, we form all the  $r$ -combinations of the set (there are  $C(n, r)$  such  $r$ -combinations).

**Then**, we generate all possible orderings in each of these  $r$ -combinations (there are  $P(r, r)$  such orderings in each case).

Therefore, we have:

$$P(n, r) = C(n, r) \cdot P(r, r)$$

Oct 1, 2015

CS 320

24

## Combinations

$$\begin{aligned}C(n, r) &= P(n, r)/P(r, r) \\ &= n!/(n-r)!/(r!(n-r)!) \\ &= n!/(r!(n-r)!)\end{aligned}$$

Now we can answer our initial question:

How many ways are there to pick a set of 3 people from a group of 6 (disregarding the order of picking)?

$$C(6, 3) = 6!/(3! \cdot 3!) = 720/(6 \cdot 6) = 720/36 = 20$$

There are 20 different ways, that is, 20 different groups that may be picked.

Oct 1, 2015

CS 320

25

## Combinations

### Corollary:

Let  $n$  and  $r$  be nonnegative integers with  $r \leq n$ . Then  $C(n, r) = C(n, n - r)$ .

Note that "picking a group of  $r$  people from a group of  $n$  people" **is the same as** "splitting a group of  $n$  people into a group of  $r$  people and another group of  $(n - r)$  people".

Please also look at the proof on page 411 (page 359 of the 6<sup>th</sup> edition).

Oct 1, 2015

CS 320

26

## Combinations

$$C(n, n-r) = \frac{n!}{(n-r)!(n-(n-r))!} = \frac{n!}{(n-r)!r!} = C(n, r)$$

This symmetry is intuitively plausible. For example, let us consider a set containing six elements ( $n = 6$ ).

**Picking two** elements and **leaving four** is essentially the same as **picking four** elements and **leaving two**.

In either case, our number of choices is the number of ways to **divide** the set into one set containing two elements and another set containing four elements.

Oct 1, 2015

CS 320

27

## Combinations

### Example:

A soccer club has 8 female and 7 male members. For today's match, the coach wants to have 6 female and 5 male players on the grass. How many possible configurations are there?

$$\begin{aligned}C(8, 6) \cdot C(7, 5) &= 8!/(6! \cdot 2!) \cdot 7!/(5! \cdot 2!) \\ &= 28 \cdot 21 \\ &= 588\end{aligned}$$

Oct 1, 2015

CS 320

28

## Combinations

### Pascal's Identity (page 418) [366 in 6<sup>th</sup> ed.]:

Let  $n$  and  $k$  be positive integers with  $n \geq k$ . Then  $C(n + 1, k) = C(n, k - 1) + C(n, k)$ .

How can this be explained?

What is it good for?

Oct 1, 2015

CS 320

29

## Combinations

Imagine a set  $S$  containing  $n$  elements and a set  $T$  containing  $(n + 1)$  elements, namely all elements in  $S$  plus a new element  $\mathbf{a}$ .

Calculating  $C(n + 1, k)$  is equivalent to answering the question: How many subsets of  $T$  containing  $k$  items are there?

Case I: The subset contains  $(k - 1)$  elements of  $S$  plus the element  $\mathbf{a}$ :  $C(n, k - 1)$  choices.

Case II: The subset contains  $k$  elements of  $S$  and does not contain  $\mathbf{a}$ :  $C(n, k)$  choices.

**Sum Rule:**  $C(n + 1, k) = C(n, k - 1) + C(n, k)$ .


Oct 1, 2015

CS 320

30


### Pascal's Triangle

In Pascal's triangle, each number is the sum of the numbers to its upper left and upper right:



### Pascal's Triangle

Since we have  $C(n + 1, k) = C(n, k - 1) + C(n, k)$  and  $C(0, 0) = 1$ , we can use Pascal's triangle to simplify the computation of  $C(n, k)$ :



### Binomial Coefficients

Expressions of the form  $C(n, k)$  are also called **binomial coefficients**.  
How come?

A **binomial expression** is the sum of two terms, such as  $(a + b)$ .  
Now consider  $(a + b)^2 = (a + b)(a + b)$ .

When expanding such expressions, we have to form all possible products of a term in the first factor and a term in the second factor:

$$(a + b)^2 = a \cdot a + a \cdot b + b \cdot a + b \cdot b$$

Then we can sum identical terms:

$$(a + b)^2 = a^2 + 2ab + b^2$$

### Binomial Coefficients

For  $(a + b)^3 = (a + b)(a + b)(a + b)$  we have  
 $(a + b)^3 = aaa + aab + aba + abb + baa + bab + bba + bbb$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

There is only one term  $a^3$ , because there is only one possibility to form it: Choose **a** from all three factors:  $C(3, 3) = 1$ .

There is three times the term  $a^2b$ , because there are three possibilities to choose **a** from a **subset** of two out of the three factors:  $C(3, 2) = 3$ .

Similarly, there is three times the term  $ab^2$  ( $C(3, 1) = 3$ ) and once the term  $b^3$  ( $C(3, 0) = 1$ ).

### Binomial Coefficients

This leads us to the following formula:

$$(a + b)^n = \sum_{j=0}^n C(n, j) \cdot a^{n-j} \cdot b^j \quad \text{(Binomial Theorem)}$$

With the help of Pascal's triangle, this formula can considerably simplify the process of expanding powers of binomial expressions.

For example, the fifth row of Pascal's triangle (1 - 4 - 6 - 4 - 1) helps us to compute  $(a + b)^4$ :

$$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$$

### Binomial Theorem

We can prove the Binomial Theorem in two ways.

First, an argument about how the term  $a^j b^{n-j}$  appears when we expand  $(a+b)^n = (a+b) \dots (a+b)$ , by choosing  $a$  from  $j$  of the factors  $(a+b)$  and  $b$  from  $n-j$  of them. This choice can be made in  $C(n, j)$  ways, so the coefficient of  $a^j b^{n-j}$  will be  $C(n, j)$

## Binomial Theorem

Binomial Theorem:  $(a+b)^n = \sum_{j=0}^n C(n,j) \cdot a^{n-j} b^j, n \geq 0$

Second proof (induction on n)

Base case:  $n = 0$ . LHS =  $(a+b)^0 = 1$

RHS =  $C(0,0)a^0b^0 = 1$ .

Induction step: Suppose the formula is true for n.

$$(a+b)^{n+1} = (a+b)(a+b)^n = (a+b) \sum_{j=0}^n C(n,j) \cdot a^{n-j} b^j$$

Oct 1, 2015

CS 320

37

$$\begin{aligned} &= \sum_{j=0}^n C(n,j) a^{n-j+1} b^j + \sum_{j=0}^n C(n,j) a^{n-j} b^{j+1} \\ &= a^{n+1} + \sum_{j=1}^n C(n,j) a^{n+1-j} b^j + \sum_{k=0}^{n-1} C(n,k) a^{(n+1)-(k+1)} b^{k+1} + b^{n+1} \\ &= a^{n+1} + \sum_{j=1}^n C(n,j) a^{n+1-j} b^j + \sum_{j=1}^n C(n,j-1) a^{(n+1)-j} b^j + b^{n+1} \\ &= a^{n+1} + \sum_{j=1}^n (C(n,j) + C(n,j-1)) a^{n+1-j} b^j + b^{n+1} \\ &= a^{n+1} + \sum_{j=1}^n C(n+1,j) a^{n+1-j} b^j + b^{n+1} = \sum_{j=0}^{n+1} C(n+1,j) a^{n+1-j} b^j \end{aligned}$$

Oct 1, 2015

CS 320

38

## Proof of Binomial Theorem

In slide 2 of the proof using induction:

From line 1 to line 2 we split off the term  $j=0$ ,  $a^n$ , from the first sum and  $k=n$ ,  $b^n$ , from the second sum (having replaced  $j$  by  $k$ ).

Note that  $C(n,0) = C(n,n) = 1$ .

From line 2 to line 3 we let  $k+1=j$ , so  $k=j-1$ , and fixed the sum limits.

Then we combined the sums and applied Pascal's theorem.

Then we put the  $a^{n+1}$ ,  $j=0$ , and  $b^{n+1}$ ,  $j=n+1$  terms back in the sum, to get the result.

Oct 1, 2015

CS 320

39