

Introduction

Modelling parallel systems

Linear Time Properties

Regular Properties

regular safety properties

ω -regular properties



model checking with Büchi automata

Linear Temporal Logic

Computation-Tree Logic

Equivalences and Abstraction

idea: define **regular LT properties** to be those languages of **infinite words** over the alphabet 2^{AP} that have a representation by a **finite automata**

idea: define **regular LT properties** to be those languages of **infinite words** over the alphabet 2^{AP} that have a representation by a **finite automata**

- regular safety properties:
NFA-representation for the **bad prefixes**

idea: define **regular LT properties** to be those languages of **infinite words** over the alphabet 2^{AP} that have a representation by a **finite automata**

- regular safety properties:
 NFA-representation for the **bad prefixes**
- representation other regular LT properties by
 - * **ω -automata**, i.e., acceptors for infinite words

idea: define **regular LT properties** to be those languages of **infinite words** over the alphabet 2^{AP} that have a representation by a **finite automata**

- regular safety properties:
 - NFA**-representation for the **bad prefixes**
- representation other regular LT properties by
 - * **ω -automata**, i.e., acceptors for infinite words
 - * **ω -regular expressions**

remind: syntax and semantics of regular expressions
over some alphabet $\Sigma = \{A, B, \dots\}$

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

where $A \in \Sigma$

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

where $A \in \Sigma$

semantics: $\alpha \mapsto \mathcal{L}(\alpha) \subseteq \Sigma^*$ language of finite words

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

where $A \in \Sigma$

semantics: $\alpha \mapsto \mathcal{L}(\alpha) \subseteq \Sigma^*$ language of finite words

$$\mathcal{L}(\emptyset) = \emptyset$$

$$\mathcal{L}(\epsilon) = \{\epsilon\}$$

$$\mathcal{L}(A) = \{A\}$$

$$\mathcal{L}(\alpha_1 + \alpha_2) = \mathcal{L}(\alpha_1) \cup \mathcal{L}(\alpha_2) \quad \text{union}$$

$$\mathcal{L}(\alpha_1 \cdot \alpha_2) = \mathcal{L}(\alpha_1)\mathcal{L}(\alpha_2) \quad \text{concatenation}$$

$$\mathcal{L}(\alpha^*) = \mathcal{L}(\alpha)^* \quad \text{Kleene closure}$$

regular expressions:

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

ω -regular expressions:

regular expressions + ω -operator α^ω

regular expressions:

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

ω -regular expressions:

regular expressions + ω -operator α^ω

Kleene star: “finite repetition”

ω -operator: “infinite repetition”

regular expressions:

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

ω -regular expressions:

regular expressions + ω -operator α^ω

Kleene star: “finite repetition”

ω -operator: “infinite repetition”

for $L \subseteq \Sigma^*$:

$$L^\omega \stackrel{\text{def}}{=} \{w_1 w_2 w_3 \dots : w_i \in L \text{ for all } i \geq 1\}$$

regular expressions:

$$\alpha ::= \emptyset \mid \epsilon \mid A \mid \alpha_1 + \alpha_2 \mid \alpha_1 \cdot \alpha_2 \mid \alpha^*$$

ω -regular expressions:

regular expressions + ω -operator α^ω

Kleene star: “finite repetition”

ω -operator: “infinite repetition”

for $L \subseteq \Sigma^*$:

$$L^\omega \stackrel{\text{def}}{=} \{w_1 w_2 w_3 \dots : w_i \in L \text{ for all } i \geq 1\}$$

note: $L^\omega \subseteq \Sigma^\omega$ if $\epsilon \notin L$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega$$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

- language of $(A^* \cdot B)^\omega$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

- language of $(A^* \cdot B)^\omega =$ set of all infinite words over $\Sigma = \{A, B\}$ containing infinitely many B 's

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

- language of $(A^* \cdot B)^\omega =$ set of all infinite words over $\Sigma = \{A, B\}$ containing infinitely many B 's
- language of $(A^* \cdot B)^\omega + (B^* \cdot A)^\omega$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

- language of $(A^* \cdot B)^\omega =$ set of all infinite words over $\Sigma = \{A, B\}$ containing infinitely many B 's
- language of $(A^* \cdot B)^\omega + (B^* \cdot A)^\omega =$ set of all infinite words over Σ with infinitely many A 's or B 's

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

- language of $(A^* \cdot B)^\omega =$ set of all infinite words over $\Sigma = \{A, B\}$ containing infinitely many B 's
- language of $(A^* \cdot B)^\omega + (B^* \cdot A)^\omega =$ set of all infinite words over Σ with infinitely many A 's or B 's $= \Sigma^\omega$

syntax of ω -regular expressions over alphabet Σ :

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega \quad \text{where}$$

α_i, β_i are regular expressions over Σ s.t. $\varepsilon \notin \mathcal{L}(\beta_i)$

semantics: the language generated by γ is:

$$\mathcal{L}_\omega(\gamma) \stackrel{\text{def}}{=} \bigcup_{1 \leq i \leq n} \mathcal{L}(\alpha_i) \mathcal{L}(\beta_i)^\omega \subseteq \Sigma^\omega$$

A language $L \subseteq \Sigma^\omega$ is called ω -regular iff there exists an ω -regular expression γ s.t.

$$L = \mathcal{L}_\omega(\gamma)$$

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

$$(A + B)^* . B^\omega$$

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

$$(A + B)^* \cdot B^\omega$$

- set of all infinite words where each A is followed immediately by letter B

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

$$(A + B)^* . B^\omega$$

- set of all infinite words where each A is followed immediately by letter B

$$(B^* . A . B)^* . B^\omega + (B^* . A . B)^\omega$$

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

$$(A + B)^* . B^\omega$$

- set of all infinite words where each A is followed immediately by letter B

$$(B^* . A . B)^* . B^\omega + (B^* . A . B)^\omega$$

- set of all infinite words where each A is followed eventually by letter B

Provide an ω -regular expression for ...

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

$$(A + B)^* . B^\omega$$

- set of all infinite words where each A is followed immediately by letter B

$$(B^* . A . B)^* . B^\omega + (B^* . A . B)^\omega$$

- set of all infinite words where each A is followed eventually by letter B

$$(B^* . A^+ . B)^* . B^\omega + (B^* . A^+ . B)^\omega$$

where $\alpha^+ \stackrel{\text{def}}{=} \alpha . \alpha^*$.

Provide an ω -regular expression for ...

alphabet $\Sigma = \{A, B\}$

- set of all infinite words over Σ containing only finitely many A 's

$$(A + B)^* . B^\omega$$

- set of all infinite words where each A is followed immediately by letter B

$$(B^* . A . B)^* . B^\omega + (B^* . A . B)^\omega$$

- set of all infinite words where each A is followed eventually by letter B

$$(B^* . A^+ . B)^* . B^\omega + (B^* . A^+ . B)^\omega \equiv (A^* . B)^\omega$$

where $\alpha^+ \stackrel{\text{def}}{=} \alpha . \alpha^*$.

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(\emptyset + \{a\} + \{a, b\})^\omega$$

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(\emptyset + \{a\} + \{a, b\})^\omega$$

Each invariant is ω -regular

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(\emptyset + \{a\} + \{a, b\})^\omega$$

Each invariant is ω -regular

Let Φ be an invariant condition and let

$$\{A \subseteq AP : A \models \Phi\} = \{A_1, \dots, A_k\}$$

Then: invariant “always Φ ” $\hat{=} (A_1 + \dots + A_k)^\omega$

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(\emptyset + \{a\} + \{a, b\})^\omega$$

Indeed: each invariant is ω -regular

- “infinitely often a ”

Let E be an LT-property over AP , i.e., $E \subseteq (2^{AP})^\omega$

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(\emptyset + \{a\} + \{a, b\})^\omega$$

Indeed: each invariant is ω -regular

- “infinitely often a ”

$$((\emptyset + \{b\})^* \cdot (\{a\} + \{a, b\}))^\omega$$

Let E be an LT-property over AP , i.e., $E \subseteq 2^{AP}$.

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$:

- “always a ” (or any other invariant)
- “infinitely often a ”
- “eventually a ”

Let E be an LT-property over AP , i.e., $E \subseteq 2^{AP}$.

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$:

- “always a ” (or any other invariant)
- “infinitely often a ”
- “eventually a ”

$$(2^{AP})^* \cdot (\{a\} + \{a, b\}) \cdot (2^{AP})^\omega$$

where $2^{AP} \hat{=} \emptyset + \{a\} + \{b\} + \{a, b\}$

Let E be an LT-property over AP , i.e., $E \subseteq 2^{AP}$.

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$:

- “always a ” (or any other invariant)
- “infinitely often a ”
- “eventually a ”

$$(2^{AP})^* \cdot (\{a\} + \{a, b\}) \cdot (2^{AP})^\omega$$

- “from some moment on a ”

Let E be an LT-property over AP , i.e., $E \subseteq 2^{AP}$.

E is called an ω -regular property iff there exists an ω -regular expression γ over 2^{AP} s.t. $E = \mathcal{L}_\omega(\gamma)$

Examples for $AP = \{a, b\}$:

- “always a ” (or any other invariant)
- “infinitely often a ”
- “eventually a ”

$$(2^{AP})^* \cdot (\{a\} + \{a, b\}) \cdot (2^{AP})^\omega$$

- “from some moment on a ”

$$(2^{AP})^* \cdot (\{a\} + \{a, b\})^\omega$$

symbolic notation for ω -regular properties

... using **formulas** instead of **sums**

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(\emptyset + \{a\} + \{a, b\})^\omega$$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(a \vee \neg b)^\omega \hat{=} (\emptyset + \{a\} + \{a, b\})^\omega$$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(a \vee \neg b)^\omega \hat{=} (\emptyset + \{a\} + \{a, b\})^\omega$$

- “infinitely often a ”

$$((\neg a)^*.a)^\omega \hat{=} ((\emptyset + \{b\})^* . (\{a\} + \{a, b\}))^\omega$$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(a \vee \neg b)^\omega \hat{=} (\emptyset + \{a\} + \{a, b\})^\omega$$

- “infinitely often a ”

$$((\neg a)^*.a)^\omega \hat{=} ((\emptyset + \{b\})^* . (\{a\} + \{a, b\}))^\omega$$

- “from some moment on a ”:

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(a \vee \neg b)^\omega \hat{=} (\emptyset + \{a\} + \{a, b\})^\omega$$

- “infinitely often a ”

$$((\neg a)^*.a)^\omega \hat{=} ((\emptyset + \{b\})^*.\{a\} + \{a, b\})^\omega$$

- “from some moment on a ”:

$$true^*.a^\omega$$

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(a \vee \neg b)^\omega \hat{=} (\emptyset + \{a\} + \{a, b\})^\omega$$

- “infinitely often a ”

$$((\neg a)^*.a)^\omega \hat{=} ((\emptyset + \{b\})^*.(\{a\} + \{a, b\}))^\omega$$

- “from some moment on a ”:

$$true^*.a^\omega$$

- “whenever a then b will hold somewhen later”

Examples for $AP = \{a, b\}$

- invariant with invariant condition $a \vee \neg b$

$$(a \vee \neg b)^\omega \hat{=} (\emptyset + \{a\} + \{a, b\})^\omega$$

- “infinitely often a ”

$$((\neg a)^*.a)^\omega \hat{=} ((\emptyset + \{b\})^* . (\{a\} + \{a, b\}))^\omega$$

- “from some moment on a ”:

$$true^*.a^\omega$$

- “whenever a then b will hold somewhen later”

$$((\neg a)^*.a.true^*.b)^* . (\neg a)^\omega + ((\neg a)^*.a.true^*.b)^\omega$$

syntax as for **NFA**



nondeterministic finite automata

syntax as for **NFA**



nondeterministic finite automata

semantics: language of infinite words

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of **final states**, also called **accept states**

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of **final states**, also called **accept states**

run for a word $A_0 A_1 A_2 \dots \in \Sigma^\omega$:

state sequence $\pi = q_0 q_1 q_2 \dots$ where $q_0 \in Q_0$
and $q_{i+1} \in \delta(q_i, A_i)$ for $i \geq 0$

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of **final states**, also called **accept states**

run for a word $A_0 A_1 A_2 \dots \in \Sigma^\omega$:

state sequence $\pi = q_0 q_1 q_2 \dots$ where $q_0 \in Q_0$
and $q_{i+1} \in \delta(q_i, A_i)$ for $i \geq 0$


run π is **accepting** if $\exists i \in \mathbb{N}. q_i \in F$

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of **final states**, also called **accept states**

accepted language $\mathcal{L}_\omega(\mathcal{A}) \subseteq \Sigma^\omega$ is given by:



$\mathcal{L}_\omega(\mathcal{A}) \stackrel{\text{def}}{=} \text{set of infinite words over } \Sigma \text{ that have an accepting run in } \mathcal{A}$



↘ initial state

○ nonfinal state

□ final state




NBA with state space $\{q_0, q_1\}$


q_0 initial state

q_1 accept state

alphabet $\Sigma = \{A, B\}$




accepted language: ?



accepted language:

set of all infinite words that contain infinitely many **A**'s




accepted language:

set of all infinite words that
contain infinitely many **A**'s

$$(B^*.A)^\omega$$

Examples for NBA over $\Sigma = \{A, B\}$


LTLMC3.2-22



accepted language:


set of all infinite words that contain infinitely many A 's

$$(B^*.A)^\omega$$



Examples for NBA over $\Sigma = \{A, B\}$


LTLMC3.2-22



accepted language:


set of all infinite words that contain infinitely many **A**'s

$$(B^*.A)^\omega$$



AABAAB**AAB**...
AAAAAAAA...


} accepted words



accepted language:

set of all infinite words that contain infinitely many **A**'s

$$(B^*.A)^\omega$$




accepted language:

“every **B** is preceded by a positive even number of **A**'s”

$AABAAABAAAB\dots$
 $AAAAAAAAAA\dots$


} accepted words



accepted language:

set of all infinite words that contain infinitely many **A**'s

$$(B^*.A)^\omega$$



accepted language:

“every **B** is preceded by a positive even number of **A**'s”

$$((A.A)^+.B)^\omega + ((A.A)^+.B)^*.A^\omega$$

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of **final states**, also called **accept states**

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$


- Q finite set of states
- Σ alphabet \leftarrow here: $\Sigma = 2^{AP}$
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of final states, also called accept states

NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

- Q finite set of states
- Σ alphabet \leftarrow here: $\Sigma = 2^{AP}$
- $\delta : Q \times \Sigma \rightarrow 2^Q$ transition relation
- $Q_0 \subseteq Q$ set of initial states
- $F \subseteq Q$ set of **final states**, also called **accept states**


accepted language $\mathcal{L}_w(\mathcal{A})$ is an **LT-property**:

$\mathcal{L}_w(\mathcal{A}) =$ set of infinite words over 2^{AP} that have an **accepting run** in \mathcal{A}




$$\mathcal{L}_\omega(\mathcal{A}) = ?$$

set of atomic propositions $AP = \{a, b\}$




$$\mathcal{L}_\omega(\mathcal{A}) \hat{=} \text{true.} \neg a. \text{true}^\omega$$


set of atomic propositions $AP = \{a, b\}$




$$\mathcal{L}_\omega(\mathcal{A}) \hat{=} \text{true} \cdot \neg a \cdot \text{true}^\omega$$



set of atomic propositions $AP = \{a, b\}$





$$\mathcal{L}_\omega(\mathcal{A}) \hat{=} \text{true} \cdot \neg a \cdot \text{true}^\omega$$




$$(a \vee \neg b) \cdot \text{true}^\omega$$

set of atomic propositions $AP = \{a, b\}$







“always a ” $\hat{=} a^\omega$




“always a ” $\hat{=} a^\omega$







“always a ” $\hat{=} a^\omega$



“infinitely often a and ...”




“always a ” $\hat{=} a^\omega$



“infinitely often a and always $a \vee b$ ”

$$\hat{=} ((a \vee b)^* . a)^\omega$$




“infinitely often a and
always $a \vee b$ ”

$$((a \vee b)^* \cdot a)^\omega$$


“infinitely often a ”

$$((\neg a)^* \cdot a)^\omega$$



“infinitely often a and
always $a \vee b$ ”

$$((a \vee b)^* . a)^\omega$$



“infinitely often a ”

$$((\neg a)^* . a)^\omega$$

For each NBA \mathcal{A} there is an ω -regular expression γ
with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

For each NBA \mathcal{A} there is an ω -regular expression γ
with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

Proof. Let \mathcal{A} be an NBA $(Q, \Sigma, \delta, Q_0, F)$

For each NBA \mathcal{A} there is an ω -regular expression γ
with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

Proof. Let \mathcal{A} be an NBA $(Q, \Sigma, \delta, Q_0, F)$ and $q, p \in Q$.
Let $\mathcal{A}_{q,p}$ be the NFA $(Q, \Sigma, \delta, q, \{p\})$.

For each NBA \mathcal{A} there is an ω -regular expression γ with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

Proof. Let \mathcal{A} be an NBA $(Q, \Sigma, \delta, Q_0, F)$ and $q, p \in Q$. Let $\mathcal{A}_{q,p}$ be the NFA $(Q, \Sigma, \delta, q, \{p\})$. Then:

$$\mathcal{L}_\omega(\mathcal{A}) = \bigcup_{q \in Q_0} \bigcup_{p \in F} \mathcal{L}(\mathcal{A}_{q,p}) (\mathcal{L}(\mathcal{A}_{p,p}) \setminus \{\varepsilon\})^\omega$$

For each NBA \mathcal{A} there is an ω -regular expression γ with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

Proof. Let \mathcal{A} be an NBA $(Q, \Sigma, \delta, Q_0, F)$ and $q, p \in Q$. Let $\mathcal{A}_{q,p}$ be the NFA $(Q, \Sigma, \delta, q, \{p\})$. Then:


$$\mathcal{L}_\omega(\mathcal{A}) = \bigcup_{q \in Q_0} \bigcup_{p \in F} \mathcal{L}(\mathcal{A}_{q,p}) (\mathcal{L}(\mathcal{A}_{p,p}) \setminus \{\varepsilon\})^\omega$$

is ω -regular as $\mathcal{L}(\mathcal{A}_{q,p})$ and $\mathcal{L}(\mathcal{A}_{p,p}) \setminus \{\varepsilon\}$ are regular


Example: NBA \rightsquigarrow ω -regular expression

LTLMC3.2-26

NBA \mathcal{A}



NBA \mathcal{A}



$$\mathcal{L}_\omega(\mathcal{A}) = L_{12}(L'_{22})^\omega \cup L_{22}(L'_{22})^\omega$$


$$L_{12} = \mathcal{L}(A_{12})$$

$$L_{22} = \mathcal{L}(A_{22})$$

$$L'_{22} = L_{22} \setminus \{\varepsilon\}$$

Example: NBA \rightsquigarrow ω -regular expression

LTLMC3.2-26


NBA \mathcal{A} 

$$\mathcal{L}_\omega(\mathcal{A}) = L_{12}(L'_{22})^\omega \cup L_{22}(L'_{22})^\omega$$

$$L_{12} = \mathcal{L}(A_{12})$$


$$L_{22} = \mathcal{L}(A_{22})$$

$$L'_{22} = L_{22} \setminus \{\varepsilon\}$$

NFA \mathcal{A}_{12} 

Example: NBA \rightsquigarrow ω -regular expression

LTLMC3.2-26

NBA \mathcal{A} 


$$\mathcal{L}_\omega(\mathcal{A}) = L_{12}(L'_{22})^\omega \cup L_{22}(L'_{22})^\omega$$

$$L_{12} = \mathcal{L}(A_{12})$$

$$L_{22} = \mathcal{L}(A_{22})$$

$$L'_{22} = L_{22} \setminus \{\varepsilon\}$$


$$L_{12} \hat{=} A.(B.A + A.A.A)^*$$

NFA \mathcal{A}_{12} 

Example: NBA \rightsquigarrow ω -regular expression

LTLMC3.2-26

NBA \mathcal{A}



$$\mathcal{L}_\omega(\mathcal{A}) = L_{12}(L'_{22})^\omega \cup L_{22}(L'_{22})^\omega$$

$$L_{12} = \mathcal{L}(A_{12})$$



$$L_{22} = \mathcal{L}(A_{22})$$

$$L'_{22} = L_{22} \setminus \{\epsilon\}$$

$$L_{12} \hat{=} A.(B.A + A.A.A)^*$$


$$L'_{22} \hat{=} (B.A + A.A.A)^+$$

NFA \mathcal{A}_{12}



Example: NBA \rightsquigarrow ω -regular expression

NBA \mathcal{A}





language of \mathcal{A} :

$$A.(B.A + A.A.A)^\omega + (B.A + A.A.A)^\omega$$

$$L_{12} \hat{=} A.(B.A + A.A.A)^*$$


$$L'_{22} \hat{=} (B.A + A.A.A)^+$$

NFA \mathcal{A}_{12}



Example: NBA \rightsquigarrow ω -regular expression

NBA \mathcal{A}





language of \mathcal{A} :

$$\begin{aligned}
 & A.(B.A + A.A.A)^\omega \\
 & + (B.A + A.A.A)^\omega \\
 \equiv & (A + \varepsilon).(B.A + A.A.A)^\omega
 \end{aligned}$$

$$L_{12} \hat{=} A.(B.A + A.A.A)^*$$

$$L'_{22} \hat{=} (B.A + A.A.A)^+$$

NFA \mathcal{A}_{12}



For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an **NBA** \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an **NBA** \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof.

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an **NBA** \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof. consider **NFA** \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an **NBA** \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof. consider **NFA** \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

- construct **NBA** \mathcal{B}_i^ω for β_i^ω

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an **NBA** \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof. consider **NFA** \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

- construct **NBA** \mathcal{B}_i^ω for β_i^ω
- construct **NBA** $\mathcal{C}_i = \mathcal{A}_i \mathcal{B}_i^\omega$ for $\alpha_i \cdot \beta_i^\omega$

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an **NBA** \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof. consider **NFA** \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

- construct **NBA** \mathcal{B}_i^ω for β_i^ω
- construct **NBA** $\mathcal{C}_i = \mathcal{A}_i \mathcal{B}_i^\omega$ for $\alpha_i \cdot \beta_i^\omega$
- construct **NBA** for $\bigcup_{1 \leq i \leq n} \mathcal{L}_\omega(\mathcal{C}_i)$

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an NBA \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof. consider NFA \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

- construct NBA \mathcal{B}_i^ω for β_i^ω
- construct NBA $\mathcal{C}_i = \mathcal{A}_i \mathcal{B}_i^\omega$ for $\alpha_i \cdot \beta_i^\omega$
- construct **NBA** for $\bigcup_{1 \leq i \leq n} \mathcal{L}_\omega(\mathcal{C}_i)$




NBA are closed under union

LTLMC3.2-28


NBA are closed under union

LTLMC3.2-28

NBA \mathcal{A}_1




NBA \mathcal{A}_2




NBA are closed under union

LTLMC3.2-28


NBA \mathcal{A}_1



NBA \mathcal{A}_2



NBA for $\mathcal{L}_w(\mathcal{A}_1) \cup \mathcal{L}_w(\mathcal{A}_2)$



For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an NBA \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.


Proof. consider NFA \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

- construct NBA \mathcal{B}_i^ω for β_i^ω
- construct **NBA** $\mathcal{C}_i = \mathcal{A}_i \mathcal{B}_i^\omega$ for $\alpha_i \cdot \beta_i^\omega$ ←
- construct NBA for $\bigcup_{1 \leq i \leq n} \mathcal{L}_\omega(\mathcal{C}_i)$


Concatenation of an NFA and an NBA

LTLMC3.2-29

NFA \mathcal{A}_1




NBA \mathcal{A}_2




Concatenation of an NFA and an NBA

LTLMC3.2-29


NFA \mathcal{A}_1



NBA \mathcal{A}_2




NBA for $\mathcal{L}(\mathcal{A}_1) \cdot \mathcal{L}_\omega(\mathcal{A}_2)$:




Concatenation of an NFA and an NBA

LTLMC3.2-29


NFA \mathcal{A}_1



NBA \mathcal{A}_2





NBA for $\mathcal{L}(\mathcal{A}_1) \cdot \mathcal{L}_\omega(\mathcal{A}_2)$:




accept states as in \mathcal{A}_2

Concatenation of an NFA and an NBA



NBA for $\mathcal{L}(\mathcal{A}_1) \cdot \mathcal{L}_\omega(\mathcal{A}_2)$:



accept states as in \mathcal{A}_2

For each ω -regular expression

$$\gamma = \alpha_1 \cdot \beta_1^\omega + \dots + \alpha_n \cdot \beta_n^\omega$$

there exists an NBA \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$.

Proof. consider NFA \mathcal{A}_i for α_i and \mathcal{B}_i for β_i

- construct NBA \mathcal{B}_i^ω for β_i^ω
- construct NBA $\mathcal{C}_i = \mathcal{A}_i \mathcal{B}_i^\omega$ for $\alpha_i \cdot \beta_i^\omega$
- construct NBA for $\bigcup_{1 \leq i \leq n} \mathcal{L}_\omega(\mathcal{C}_i)$




ω -operator for NFA

LTLMC3.2-30

NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$




NBA \mathcal{A}^ω for language
 $L^\omega \subseteq \Sigma^\omega$



ω -operator for NFA


LTLMC3.2-30

NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



\rightsquigarrow


NBA \mathcal{A}^ω for language
 $L^\omega \subseteq \Sigma^\omega$



ω -operator for NFA


LTLMC3.2-30

NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



\rightsquigarrow

NBA \mathcal{A}^ω for language
 $L^\omega \subseteq \Sigma^\omega$




wrong !

ω -operator for NFA


LTLMC3.2-30

NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



\rightsquigarrow

NBA \mathcal{A}^ω for language
 $L^\omega \subseteq \Sigma^\omega$




wrong !


ω -operator for NFA

LTLMC3.2-30

NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$

 \rightsquigarrow


NBA \mathcal{A}^ω for language
 $L^\omega \subseteq \Sigma^\omega$





wrong !

... correct, if $\delta(q, x) = \emptyset \quad \forall q \in F \quad \forall x \in \Sigma$

NFA \mathcal{A} for language $L \subseteq \Sigma^+$ \implies NFA \mathcal{B} for L s.t. all final states are terminal



NFA \mathcal{A} for language $L \subseteq \Sigma^+$ \implies NFA \mathcal{B} for L s.t. all final states are terminal



... add a new final state p' ...

ω -operator for NFA

LTLMC3.2-31



NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



NFA \mathcal{B} for L s.t. all
final states are terminal



NBA \mathcal{B}^ω



... add a new final state p' ...

ω -operator for NFA

LTLMC3.2-31



NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



NFA \mathcal{B} for L s.t. all
final states are terminal



NBA \mathcal{B}^ω



... add a new final state p' ...

ω -operator for NFA

LTLMC3.2-31



NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



NFA \mathcal{B} for L s.t. all
final states are terminal



NBA \mathcal{B}^ω



... add a new final state p' ...

ω -operator for NFA

LTLMC3.2-31



NFA \mathcal{A} for language
 $L \subseteq \Sigma^+$



NFA \mathcal{B} for L s.t. all
final states are terminal




NBA \mathcal{B}^ω



$$\mathcal{L}(\mathcal{A})^\omega = \mathcal{L}_\omega(\mathcal{B}^\omega)$$



Example: ω -operator for NFA

LTLMC3.2-32






Example: ω -operator for NFA

LTLMC3.2-32






Example: ω -operator for NFA

LTLMC3.2-32



Example: ω -operator for NFA

LTLMC3.2-32



- (1) For each NBA \mathcal{A} there exists an ω -regular expression γ with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$
- (2) For each ω -regular expression γ there exists an NBA \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

- (1) For each NBA \mathcal{A} there exists an ω -regular expression γ with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$
- (2) For each ω -regular expression γ there exists an NBA \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

Corollary:

If E be an LT property then:

E is ω -regular iff $E = \mathcal{L}_\omega(\mathcal{A})$ for some NBA \mathcal{A}

- (1) For each NBA \mathcal{A} there exists an ω -regular expression γ with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$
- (2) For each ω -regular expression γ there exists an NBA \mathcal{A} with $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\gamma)$

Corollary:

If E be an LT property, i.e., $E \subseteq (2^{AP})^\omega$, then:

E is ω -regular iff $E = \mathcal{L}_\omega(\mathcal{A})$ for some NBA \mathcal{A} over the alphabet 2^{AP}

remind: Kleene's theorem for regular languages:

The class of **regular languages** is closed under

- union, intersection, complementation
- concatenation and Kleene star

remind: Kleene's theorem for regular languages:

The class of **regular languages** is closed under

- **union, intersection, complementation**
- concatenation and Kleene star

The class of **ω -regular languages** is closed under **union, intersection** and **complementation**.

The class of ω -regular languages is closed under union, intersection and complementation.

- *union:*
- *intersection:*
- *complementation:*

The class of ω -regular languages is closed under union, intersection and complementation.

- *union*:
obvious from definition of ω -regular expressions
- *intersection*:

- *complementation*:

The class of ω -regular languages is closed under union, intersection and complementation.

- *union*:
obvious from definition of ω -regular expressions
- *intersection*:
will be discussed later
relies on a certain product construction for NBA
- *complementation*:

The class of ω -regular languages is closed under union, intersection and complementation.

- *union*:
obvious from definition of ω -regular expressions
- *intersection*:
will be discussed later
relies on a certain product construction for NBA
- *complementation*:
much more difficult than for NFA,
via other types of ω -automata

Nonemptiness for NBA

LTLMC3.2-NBA-EMPTINESS

given: NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$

question: does $\mathcal{L}_\omega(\mathcal{A}) \neq \emptyset$ hold ?

Let $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ be an NBA. Then:

$$\mathcal{L}_\omega(\mathcal{A}) \neq \emptyset \quad \text{iff} \quad \exists q_0 \in Q_0 \exists p \in F \exists x \in \Sigma^* \exists y \in \Sigma^+.$$
$$p \in \delta(q_0, x) \cap \delta(p, y)$$


Nonemptiness for NBA

Let $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ be an NBA. Then:

$$\mathcal{L}_\omega(\mathcal{A}) \neq \emptyset \quad \text{iff} \quad \exists q_0 \in Q_0 \exists p \in F \exists x \in \Sigma^* \exists y \in \Sigma^+.$$
$$p \in \delta(q_0, x) \cap \delta(p, y)$$




there exists a reachable accept state $p \in F$
that belongs to a cycle



Nonemptiness for NBA

Let $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ be an NBA. Then:

$$\begin{aligned} \mathcal{L}_\omega(\mathcal{A}) \neq \emptyset & \text{ iff } \exists q_0 \in Q_0 \exists p \in F \exists x \in \Sigma^* \exists y \in \Sigma^+. \\ & p \in \delta(q_0, x) \cap \delta(p, y) \\ & \text{ iff there exist finite words } x, y \in \Sigma^* \\ & \text{ s.t. } y \neq \varepsilon \text{ and } xy^\omega \in \mathcal{L}_\omega(\mathcal{A}) \end{aligned}$$




Nonemptiness for NBA

Let $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ be an NBA. Then:

$$\mathcal{L}_\omega(\mathcal{A}) \neq \emptyset \quad \text{iff} \quad \exists q_0 \in Q_0 \exists p \in F \exists x \in \Sigma^* \exists y \in \Sigma^+.$$
$$p \in \delta(q_0, x) \cap \delta(p, y)$$

iff there exist finite words $x, y \in \Sigma^*$
s.t. $y \neq \varepsilon$ and $xy^\omega \in \mathcal{L}_\omega(\mathcal{A})$

↑
“ultimately periodic words”



Let $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ be an NBA. Then:

$$\begin{aligned} \mathcal{L}_\omega(\mathcal{A}) \neq \emptyset & \text{ iff } \exists q_0 \in Q_0 \exists p \in F \exists x \in \Sigma^* \exists y \in \Sigma^+. \\ & \quad p \in \delta(q_0, x) \cap \delta(p, y) \\ & \text{ iff } \text{there exist finite words } x, y \in \Sigma^* \\ & \quad \text{s.t. } y \neq \varepsilon \text{ and } xy^\omega \in \mathcal{L}_\omega(\mathcal{A}) \end{aligned}$$

The emptiness problem for NBA is solvable by means of graph algorithms in time $\mathcal{O}(\text{poly}(\mathcal{A}))$

A DBA is an NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ such that

- \mathcal{A} has a unique initial state,
- $|\delta(q, A)| \leq 1$ for all $q \in Q$ and $A \in \Sigma$

A DBA is an NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ such that

- \mathcal{A} has a unique initial state, i.e., Q_0 is a singleton
- $|\delta(q, A)| \leq 1$ for all $q \in Q$ and $A \in \Sigma$

A DBA is an NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ such that


- \mathcal{A} has a unique initial state, i.e., Q_0 is a singleton
- $|\delta(q, A)| \leq 1$ for all $q \in Q$ and $A \in \Sigma$

notation: $\mathcal{A} = (Q, \Sigma, \delta, q_0, F)$ if $Q_0 = \{q_0\}$

A DBA is an NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ such that

- \mathcal{A} has a unique initial state, i.e., Q_0 is a singleton
- $|\delta(q, A)| \leq 1$ for all $q \in Q$ and $A \in \Sigma$

notation: $\mathcal{A} = (Q, \Sigma, \delta, q_0, F)$ if $Q_0 = \{q_0\}$




alphabet $\Sigma = \{A, B\}$

A DBA is an NBA $\mathcal{A} = (Q, \Sigma, \delta, Q_0, F)$ such that

- \mathcal{A} has a unique initial state, i.e., Q_0 is a singleton
- $|\delta(q, A)| \leq 1$ for all $q \in Q$ and $A \in \Sigma$

notation: $\mathcal{A} = (Q, \Sigma, \delta, q_0, F)$ if $Q_0 = \{q_0\}$



DBA for “infinitely often B ”

alphabet $\Sigma = \{A, B\}$

well-known:

the **powerset construction** for the
determinization (and complementation) of
finite automata (NFA)


well-known:

the powerset construction for the
determinization (and complementation) of
finite automata (NFA)


question:

does the powerset construction also work for
Büchi automata (NBA) ?


NBA for “eventually forever a ”




NBA for “eventually forever a ”




NBA for “eventually forever a ”




powerset construction



NBA for “eventually forever a ”




powerset construction




e.g., $\delta(q_0, a) = \{q_0, q_F\}$ and $\delta(q_0, \neg a) = \{q_0\}$

NBA for “eventually forever a ”




powerset construction




e.g., $\delta(q_0, a) = \{q_0, q_F\}$ and $\delta(q_0, \neg a) = \{q_0\}$

NBA for “eventually forever a ”




powerset construction




DBA for “infinitely often a ”

NBA for “eventually forever a ”



powerset construction




DBA for “infinitely often a ”

well-known:


DFA can be complemented by
complementation of the acceptance set

question:

does this also work for **DBA** ?





DBA for
“infinitely often $\neg a$ ”



DBA for
“infinitely often $\neg a$ ”


complement automaton






DBA for
“infinitely often $\neg a$ ”

complement automaton




DBA for
“infinitely often a ”




DBA for
“infinitely often $\neg a$ ”

complement automaton




DBA for
“infinitely often a ”



DBA for
“infinitely often $\neg a$ ”

complement automaton



DBA for
“infinitely often a ”

There is **no DBA** for the LT-property
“eventually forever a ”

There is no DBA \mathcal{A} over the alphabet $\Sigma = \{A, B\}$
such that $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega((A + B)^* \cdot A^\omega)$

There is no DBA \mathcal{A} over the alphabet $\Sigma = \{A, B\}$
such that $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega((A + B)^* \cdot A^\omega)$

Hence: there is no DBA for the LT-property
“eventually forever a ”

There is no DBA \mathcal{A} over the alphabet $\Sigma = \{A, B\}$
such that $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega((A + B)^* \cdot A^\omega)$

Hence: there is no DBA for the LT-property
“eventually forever a ”

Proof: apply the above theorem for $A = \{a\}$, $B = \emptyset$

There is no DBA \mathcal{A} over the alphabet $\Sigma = \{A, B\}$ such that $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega((A + B)^* \cdot A^\omega)$

Hence: there is no DBA for the LT-property
“eventually forever a ”

Proof: apply the above theorem for $A = \{a\}$, $B = \emptyset$

The class of **DBA-recognizable languages** is a proper subclass of the class of ω -regular languages

There is no DBA \mathcal{A} over the alphabet $\Sigma = \{A, B\}$ such that $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega((A + B)^* \cdot A^\omega)$

Hence: there is no DBA for the LT-property
“eventually forever a ”

Proof: apply the above theorem for $A = \{a\}$, $B = \emptyset$

The class of **DBA-recognizable languages** is a proper subclass of the class of ω -regular languages and is not closed under complementation.

There is no DBA \mathcal{A} over the alphabet $\Sigma = \{A, B\}$ such that $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega((A + B)^* \cdot A^\omega)$

The class of **DBA-recognizable languages** is a proper subclass of the class of ω -regular languages and is not closed under complementation.

$(A^* \cdot B)^\omega$ “infinitely many B 's” DBA-recognizable

$(A + B)^* \cdot A^\omega$ “only finitely many B 's”
not DBA-recognizable

A generalized nondeterministic Büchi automaton is a tuple

$$\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$$

where Q, Σ, δ, Q_0 are as in NBA, but \mathcal{F} is a set of **accept sets**, i.e., $\mathcal{F} \subseteq 2^Q$.

A generalized nondeterministic Büchi automaton is a tuple

$$\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$$

where Q, Σ, δ, Q_0 are as in NBA, but \mathcal{F} is a set of **accept sets**, i.e., $\mathcal{F} \subseteq 2^Q$.

A run $q_0 q_1 q_2 \dots$ for some infinite word $\sigma \in \Sigma^\omega$ is called **accepting** if **each accept set** is visited infinitely often

A generalized nondeterministic Büchi automaton is a tuple

$$\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$$

where Q, Σ, δ, Q_0 are as in NBA, but \mathcal{F} is a set of **accept sets**, i.e., $\mathcal{F} \subseteq 2^Q$.

A run $q_0 q_1 q_2 \dots$ for some infinite word $\sigma \in \Sigma^\omega$ is called **accepting** if **each accept set** is visited infinitely often, i.e.,

$$\forall F \in \mathcal{F} \exists^\infty i \in \mathbb{N} \text{ s.t. } q_i \in F$$

GNBA $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ as NBA, but $\mathcal{F} \subseteq 2^Q$


A run $q_0 q_1 q_2 \dots$ for some infinite word $\sigma \in \Sigma^\omega$ is accepting if

$$\forall F \in \mathcal{F} \exists i \in \mathbb{N} \text{ s.t. } q_i \in F$$

accepted language:


$$\mathcal{L}_\omega(\mathcal{G}) \stackrel{\text{def}}{=} \{ \sigma \in \Sigma^\omega : \sigma \text{ has an accepting run in } \mathcal{G} \}$$

GNBA \mathcal{G} over $\Sigma = 2^{AP}$ where $AP = \{\text{crit}_1, \text{crit}_2\}$



$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

GNBA \mathcal{G} over $\Sigma = 2^{AP}$ where $AP = \{\text{crit}_1, \text{crit}_2\}$




$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

specifies the LT-property

“infinitely often **crit₁** and infinitely often **crit₂**”

GNBA \mathcal{G} over $\Sigma = 2^{AP}$ where $AP = \{\text{crit}_1, \text{crit}_2\}$




$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

note: $q_0 \xrightarrow{A} q_1$ implies $A \models \text{crit}_1$

$q_0 \xrightarrow{A} q_2$ implies $A \models \text{crit}_2$

GNBA \mathcal{G} over $\Sigma = 2^{AP}$ where $AP = \{\text{crit}_1, \text{crit}_2\}$



$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$


note: $q_0 \xrightarrow{A} q_1$ implies $A \models \text{crit}_1$

$q_0 \xrightarrow{A} q_2$ implies $A \models \text{crit}_2$

hence: if $A_0 A_1 A_2 \dots \in \mathcal{L}_\omega(\mathcal{G})$ then

$$\exists^\infty i \geq 0. \text{crit}_1 \in A_i \wedge \exists^\infty i \geq 0. \text{crit}_2 \in A_i$$

GNBA \mathcal{G} over $\Sigma = 2^{AP}$ where $AP = \{\text{crit}_1, \text{crit}_2\}$




$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

all words $A_0 A_1 A_2 \dots \in \Sigma^\omega$ s.t. $\exists i \geq 0. \text{crit}_1 \in A_i$ and $\exists i \geq 0. \text{crit}_2 \in A_i$ have an accepting run of the form:


$q_0 \dots q_0 q_1 q_0 \dots q_0 q_2 q_0 \dots q_0 q_1 q_0 \dots q_0 q_2 \dots$

GNBA \mathcal{G}



$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$


GNBA \mathcal{G}



$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

$$\mathcal{L}_\omega(\mathcal{G}) = ?$$

GNBA \mathcal{G}




$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

$$\mathcal{L}_\omega(\mathcal{G}) = \emptyset$$


Examples: GNBA over $\Sigma = \{A, B\}$

LTLMC3.2-41

GNBA \mathcal{G} 


$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

$$\mathcal{L}_\omega(\mathcal{G}) = \emptyset$$

GNBA \mathcal{G}' with $\mathcal{F}' = \{\{q_1, q_3\}, \{q_2, q_4\}\}$ 


Examples: GNBA over $\Sigma = \{A, B\}$

LTLMC3.2-41

GNBA \mathcal{G} 


$$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$$

$$\mathcal{L}_\omega(\mathcal{G}) = \emptyset$$

GNBA \mathcal{G}' with $\mathcal{F}' = \{\{q_1, q_3\}, \{q_2, q_4\}\}$ 

accepted language: ?


GNBA \mathcal{G}



$\mathcal{F} = \{\{q_1\}, \{q_2\}\}$

$\mathcal{L}_\omega(\mathcal{G}) = \emptyset$

GNBA \mathcal{G}' with $\mathcal{F}' = \{\{q_1, q_3\}, \{q_2, q_4\}\}$




accepted language: $A.B^\omega + A.B^+.A.(A.B)^\omega$

Empty acceptance condition


LTLMC3.2-42

NBA \mathcal{A} over $\Sigma = \{A, B\}$:



acceptance set $F = \emptyset$

GNBA \mathcal{G} over $\Sigma = \{A, B\}$:




set of acceptance sets
 $\mathcal{F} = \emptyset$

Empty acceptance condition

LTLMC3.2-42


NBA \mathcal{A} over $\Sigma = \{A, B\}$:



acceptance set $F = \emptyset$

$$\mathcal{L}_\omega(\mathcal{A}) = \emptyset$$

GNBA \mathcal{G} over $\Sigma = \{A, B\}$:




set of acceptance sets

$$\mathcal{F} = \emptyset$$

Empty acceptance condition

LTLMC3.2-42


NBA \mathcal{A} over $\Sigma = \{A, B\}$:



acceptance set $F = \emptyset$

$$\mathcal{L}_\omega(\mathcal{A}) = \emptyset$$

GNBA \mathcal{G} over $\Sigma = \{A, B\}$:



set of acceptance sets


$$\mathcal{F} = \emptyset$$

$$\mathcal{L}_\omega(\mathcal{G}) = ?$$

Empty acceptance condition

LTLMC3.2-42


NBA \mathcal{A} over $\Sigma = \{A, B\}$:



acceptance set $F = \emptyset$

$$\mathcal{L}_\omega(\mathcal{A}) = \emptyset$$

GNBA \mathcal{G} over $\Sigma = \{A, B\}$:



set of acceptance sets


$$\mathcal{F} = \emptyset$$

$$\mathcal{L}_\omega(\mathcal{G}) = \{A^\omega\}$$

Empty acceptance condition

LTLMC3.2-42


NBA \mathcal{A} over $\Sigma = \{A, B\}$:



acceptance set $F = \emptyset$

$$\mathcal{L}_\omega(\mathcal{A}) = \emptyset$$

GNBA \mathcal{G} over $\Sigma = \{A, B\}$:



set of acceptance sets

$$\mathcal{F} = \emptyset$$

$$\mathcal{L}_\omega(\mathcal{G}) = \{A^\omega\}$$

$$\mathcal{L}_\omega(\mathcal{G}) = \left\{ \begin{array}{l} \text{set of all infinite words} \\ \text{that have an infinite run} \end{array} \right.$$

For every GNBA \mathcal{G} there exists a GNBA \mathcal{G}' such that

- $\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{G}')$
- the set of acceptance sets of \mathcal{G}' is nonempty

For every GNBA \mathcal{G} there exists a GNBA \mathcal{G}' such that

- $\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{G}')$
- the set of acceptance sets of \mathcal{G}' is nonempty

correct

$$\text{GNBA } \mathcal{G} = (Q, \Sigma, \delta, Q_0, \emptyset)$$



$$\text{GNBA } \mathcal{G}' = (Q, \Sigma, \delta, Q_0, \{Q\})$$

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$

Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$

Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$
and $k \geq 1$

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$

Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$
and $k \geq 1$

note: if $k = 1$ then \mathcal{G} is an NBA

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$

Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$
and $k \geq 2$

note: if $k = 1$ then \mathcal{G} is an NBA

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with


$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$

Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$ and $k \geq 2$. NBA \mathcal{A} results from k copies of \mathcal{G} :

For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$


Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$ and $k \geq 2$. NBA \mathcal{A} results from k copies of \mathcal{G} :



For each **GNBA** \mathcal{G} there exists an **NBA** \mathcal{A} with

$$\mathcal{L}_\omega(\mathcal{G}) = \mathcal{L}_\omega(\mathcal{A})$$


Proof. Let $\mathcal{G} = (Q, \Sigma, \delta, Q_0, \mathcal{F})$ with $\mathcal{F} = \{F_1, \dots, F_k\}$ and $k \geq 2$. NBA \mathcal{A} results from k copies of \mathcal{G} :



size of the NBA: $size(\mathcal{A}) = \mathcal{O}(size(\mathcal{G}) \cdot |\mathcal{F}|)$

Example: from GNBA to NBA

LTLMC3.2-45




alphabet $\Sigma = 2^{AP}$ where
 $AP = \{\text{crit}_1, \text{crit}_2\}$

infinitely often crit_1 and
infinitely often crit_2


Example: from GNBA to NBA

LTLMC3.2-45



alphabet $\Sigma = 2^{AP}$ where
 $AP = \{crit_1, crit_2\}$

infinitely often $crit_1$ and
infinitely often $crit_2$



NBA \mathcal{A}

The class of ω -regular languages is closed under union, intersection and complementation.

The class of ω -regular languages is closed under union, intersection and complementation.

- *union*:
obvious from definition of ω -regular expressions
- *intersection*:
via some product construction
- *complementation*:
via other types of ω -automata
(not discussed here)

The class of ω -regular languages is closed under union, intersection and complementation.

- *union*:
obvious from definition of ω -regular expressions
- *intersection*: ← using **GNBA**
via some product construction
- *complementation*:
via other types of ω -automata
(not discussed here)

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_w(\mathcal{A}) = \mathcal{L}_w(\mathcal{A}_1) \cap \mathcal{L}_w(\mathcal{A}_2)$

recall:

intersection for finite automata NFA \mathcal{A}_1 and \mathcal{A}_2 is realized by a **product construction** that

- runs \mathcal{A}_1 and \mathcal{A}_2 in parallel (synchronously)
- checks whether both end in a final state

$$\left. \begin{array}{l} \mathcal{A}_1 = (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 = (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{array} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

idea: define $\mathcal{A}_1 \otimes \mathcal{A}_2$ as for NFA, i.e.,

- \mathcal{A}_1 and \mathcal{A}_2 run in parallel (synchronously)
- and check whether both are accepting

$$\left. \begin{array}{l} \mathcal{A}_1 = (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 = (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{array} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

idea: define $\mathcal{A}_1 \otimes \mathcal{A}_2$ as for NFA, i.e.,

- \mathcal{A}_1 and \mathcal{A}_2 run in parallel (synchronously)
- and check whether both are accepting



i.e., both F_1 and F_2 are visited infinitely often

$$\left. \begin{array}{l} \mathcal{A}_1 = (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 = (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{array} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

idea: define $\mathcal{A}_1 \otimes \mathcal{A}_2$ as for NFA, i.e.,

- \mathcal{A}_1 and \mathcal{A}_2 run in parallel (synchronously)
- and check whether both are accepting



i.e., both F_1 and F_2 are visited infinitely often

\rightsquigarrow product of \mathcal{A}_1 and \mathcal{A}_2 yields a **GNBA**

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\text{GNBA } \mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2$$

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\text{GNBA } \mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2$$

- state space $Q = Q_1 \times Q_2$

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\text{GNBA } \mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2$$

- state space $Q = Q_1 \times Q_2$
- alphabet Σ

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\text{GNBA } \mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2$$

- state space $Q = Q_1 \times Q_2$
- alphabet Σ
- set of initial states: $Q_0 = Q_{0,1} \times Q_{0,2}$

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\text{GNBA } \mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2$$

- state space $Q = Q_1 \times Q_2$
- alphabet Σ
- set of initial states: $Q_0 = Q_{0,1} \times Q_{0,2}$
- acceptance condition: $\mathcal{F} = \{F_1 \times Q_2, Q_1 \times F_2\}$

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

$$\text{GNBA } \mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2$$

- state space $Q = Q_1 \times Q_2$
- alphabet Σ
- set of initial states: $Q_0 = Q_{0,1} \times Q_{0,2}$
- acceptance condition: $\mathcal{F} = \{F_1 \times Q_2, Q_1 \times F_2\}$
- transition relation:

$$\delta(\langle q_1, q_2 \rangle, A) = \{ \langle p_1, p_2 \rangle : p_1 \in \delta_1(q_1, A), p_2 \in \delta_2(q_2, A) \}$$

$$\left. \begin{aligned} \mathcal{A}_1 &= (Q_1, \Sigma, \delta_1, Q_{0,1}, F_1) \\ \mathcal{A}_2 &= (Q_2, \Sigma, \delta_2, Q_{0,2}, F_2) \end{aligned} \right\} \text{two NBA}$$

goal: define an NBA \mathcal{A} s.t. $\mathcal{L}_\omega(\mathcal{A}) = \mathcal{L}_\omega(\mathcal{A}_1) \cap \mathcal{L}_\omega(\mathcal{A}_2)$

GNBA $\mathcal{G} = \mathcal{A}_1 \otimes \mathcal{A}_2 \rightsquigarrow$ equivalent NBA \mathcal{A}

- state space $Q = Q_1 \times Q_2$
- alphabet Σ
- set of initial states: $Q_0 = Q_{0,1} \times Q_{0,2}$
- acceptance condition: $\mathcal{F} = \{F_1 \times Q_2, Q_1 \times F_2\}$
- transition relation:

$$\delta(\langle q_1, q_2 \rangle, A) = \{ \langle p_1, p_2 \rangle : p_1 \in \delta_1(q_1, A), p_2 \in \delta_2(q_2, A) \}$$

Summary: ω -regular languages

LTLMC3.2-45C

The class of ω -regular languages agrees with

- the class of languages given by ω -regular expressions
- the class of **NBA**-recognizable languages
- the class of **GNBA**-recognizable languages

The class of ω -regular languages agrees with

- the class of languages given by ω -regular expressions
- the class of **NBA**-recognizable languages
- the class of **GNBA**-recognizable languages

but **DBA** are strictly less expressive

The class of ω -regular languages agrees with

- the class of languages given by ω -regular expressions
- the class of **NBA**-recognizable languages
- the class of **GNBA**-recognizable languages

but **DBA** are strictly less expressive

The class of ω -regular languages is closed under union, intersection and complementation.