# Every Incomplete Computably Enumerable Truth-Table Degree Is Branching*

Peter A. Fejer
University of Massachusetts at Boston

Richard A. Shore**
Cornell University

January 20, 1999

**Abstract**

If $r$ is a reducibility between sets of numbers, a natural question to ask about the structure $\mathcal{C}_r$ of the $r$-degrees containing computably enumerable sets is whether every element not equal to the greatest one is branching (i.e., the meet of two elements strictly above it). For the commonly studied reducibilities, the answer to this question is known except for the case of truth-table (tt) reducibility. In this paper, we answer the question in the tt case by showing that every tt-incomplete computably enumerable truth-table degree $\mathbf{a}$ is branching in $\mathcal{C}_{tt}$. The fact that every Turing-incomplete computably enumerable truth-table degree is branching can be shown using a technique of Ambos-Spies. Although this result has been known for some time, it has not appeared in the literature, and we give a proof here. The proof uses an infinite injury argument. Our main result is the proof when $\mathbf{a}$ is Turing-complete but tt-incomplete. Here we are able to exploit the Turing-completeness of $\mathbf{a}$ in a novel way to give a finite injury proof.

## 1 Introduction

If $r$ is one of the reducibilities between sets of natural numbers studied in computability theory, one can form the structures $\mathcal{D}_r$, $\mathcal{D}_r(\leq \mathbf{0}'_r)$ and $\mathcal{C}_r$ consisting of all the $r$-degrees, the $r$-degrees of sets $r$-reducible to the 1-complete computably enumerable set $K = \{e|\{e\}(e)\downarrow\}$, and the $r$-degrees containing computably enumerable sets, respectively. (We write $\mathbf{0}'_r$ for the $r$-degree of $K$.) These structures are always partially ordered sets and are usually upper semi-lattices. Among the many algebraic questions one might ask about these structures, a basic one

*The authors thank André Nies for helpful conversations.

**Partially supported by NSF Grants DMS-9503503 and DMS-9802843 and as a Visiting Scholar by M.I.T. and Harvard University.

is whether every element not equal to the greatest element is branching, where we call a degree *branching* if it is meet-reducible, i.e., it is the meet of two degrees strictly above it. In the case of $\mathcal{C}_r$ and the most commonly studied reducibilities $r$ (namely, many-one (m), truth-table (tt), weak truth-table (wtt) and Turing (T)) we have the following results. Every incomplete c.e. $m$-degree is branching. This can be seen as follows. Given a c.e. $m$-degree, $\mathbf{c} < \mathbf{0}'_m$, by a result of Denisov [2], there is a c.e. $m$-degree $\mathbf{a}$ with $\mathbf{c} < \mathbf{a} < \mathbf{0}'_m$. Now, by a result of Ershov and Lavrov [3], there is a strong minimal cover $\mathbf{b}$ of $\mathbf{c}$ in the c.e. $m$-degrees (that is, $\mathbf{c} < \mathbf{b}$ and for all c.e. $m$-degrees $\mathbf{d}$, if $\mathbf{d} < \mathbf{b}$, then $\mathbf{d} \leq \mathbf{c}$) such that $\mathbf{b}$ is incomparable with $\mathbf{a}$. The two degrees $\mathbf{a}$ and $\mathbf{b}$ now have meet $\mathbf{c}$. (See Odifreddi [8], Exercise X.5.5 for a direct proof of the fact that every c.e. $m$-degree is branching.) Every incomplete c.e. wtt-degree is branching by a result of Cohen [1]. (See Odifreddi [8], Exercise X.6.8.) In the case of Turing reducibility, the situation is a bit more complicated. There are nonzero branching c.e. T-degrees and $\mathbf{0}$ is also branching, but there are also incomplete c.e. T-degrees that are not branching (Lachlan [6]). In fact, both the branching (Slaman [9]) and nonbranching (Fejer [4]) c.e. T-degrees are dense in $\mathcal{C}_T$. Thus, among these problems, only the truth-table case has been unresolved. In this paper, we show that every incomplete c.e. truth-table degree is branching.

An unpublished argument using a technique due to Ambos-Spies shows that every Turing-incomplete c.e. truth-table degree is branching. (We call a c.e. truth-table degree *Turing-complete* if its members belong to $deg_T(K)$.) Ambos-Spies' technique was used by Nies and Shore [7] in showing a related result. We begin this paper by showing how the result in [7] can be strengthened slightly in order to get a result that has as a corollary the fact that every Turing-incomplete c.e. truth-table degree is branching. Then, in the main part of the paper, we show (Theorem 4) that every Turing-complete, tt-incomplete truth-table degree is branching. In combination with the result on Turing-incomplete c.e. degrees, this gives the result of the title of this paper. We conclude the paper by mentioning some related open problems.

Our notation is for the most part standard. (See for example Soare[10].) Our notation for truth-table reductions is from [5]. In particular, if $\{e\}(x)\downarrow$, then $[e](x)$ denotes the truth-table with Gödel number $\{e\}(x)$ and $|[e](x)|$ denotes the length of this truth table. (Here, as in [5], we take this length to be one more than the largest number asked about in the truth table.) For any set $A$, $[e]^A(x)$ is 1 or 0 according to whether or not $A$ satisfies $[e](x)$. We then have that for any two sets $A$ and $B$, $A \leq_{tt} B$ if and only if there is an $e$ with $\{e\}$ total such that for all $x$, $[e]^B(x) = A(x)$. Also, we let $F_k$ denote the finite set with canonical index $k$.

## 2 Turing Incomplete Truth-Table Degrees

In this section, we present a slight strengthening of a result from [7] which will allow us to conclude that every Turing incomplete c.e. truth-table degree is branching. We first give a slightly modified version of Lemma 3.3 of [7].

**Lemma 1** *Let $(D_n)_{n\in\omega}$ be a uniformly c.e. sequence of sets and let*

$$D = \{\langle x, n\rangle | x \in D_n\}.$$

*Suppose that, for each $e$, the set $D^{[\leq e]}$ is $T$-incomplete. Then, there exist c.e. sets $B, C$ such that*

$$(\forall n)[D_n \leq_{tt} B, C] \tag{1}$$

$$(\forall n)[B, C \not\leq_{tt} D_n] \tag{2}$$

$$(\forall Z)[Z \leq_{tt} B, C \Rightarrow (\exists e)[Z \leq_{tt} D^{[\leq e]}]] \tag{3}$$

*We say that $B, C$ form an exact pair for the sequence $(D_n)$.*

**Proof:** The difference between this result and Lemma 3.3 of [7] is the addition of equation (2). We will not give a complete proof here, but instead will just describe the modifications that need to be made to the proof of Lemma 3.3 of [7]. We have the requirements $P_n$ and $Q_e$ of [7]. To make (2) hold, we introduce requirements

$$R^X_{e,n}: \quad X \neq [e]^{D^{[n]}} \quad (X = B, C)$$

for $e, n \in \omega$. We let $(R_i)_{i\in\omega}$ be an effective listing of the $R^X_{e,n}$. For each $\gamma$ in the tree $\omega^{<\omega}$, there are strategies $P_\gamma, Q_\gamma, R_\gamma$ for $P_i, Q_i, R_i$, respectively, where $i = |\gamma|$. Strategies $P_\gamma$ and $Q_\gamma$ work as in [7], except that $P_\gamma$ tries to code $D^{[i]}$ into $X^{[2n(\gamma)]}$ for $X = B, C$, instead of into $X^{[n(\gamma)]}$, where $n(\gamma)$ is the code number assigned to $\gamma$.

The strategy for $R_\gamma$ is Sacks coding; that is, if $R_{|\gamma|}$ is $R^X_{e,n}$, then $R_\gamma$ codes $K$ into $X$ (more exactly into $X^{[2n(\gamma)+1]}$) until a difference between $X$ and $[e]^{D^{[n]}}$ occurs. (This must happen since $D^{[n]}$ is tt-incomplete.) Because the reduction being diagonalized against is a truth-table reduction, the action of $R_\gamma$ is finite.

To measure the length of agreement for the $R_i$ requirements, we define

$$l'(i, s) = \max\{x | (\forall y < x)[X(y) = [e]^{D^{[n]}}(y)[s]]\},$$

where $R_i = R^X_{e,n}$. The construction is the same as in [7] except for the following changes. 1) In Step 1, we consider numbers $x = \langle z, 2n(\beta)\rangle$ instead of $x = \langle z, n(\beta)\rangle$. 2) We add a Step 1.5 (where $R_\beta$ codes $K$) given by

For each $x < s$, if $l'(e,s) \geq x \geq R(\beta,[s,e])$ and $x = \langle z, 2n(\beta)+1\rangle$ for $z \in K_s$, then enumerate $x$ into $X$, where $R_e$ is $R^X_{i,n}$.

3) In Step 3, replace equation (5.9) by

$$F \cap (\omega^{[2n(\gamma)]} \cup \omega^{[2n(\gamma)+1]}) = \emptyset \text{ for each } \gamma < \beta * m.$$

In the verification, Lemma 5.1 of [7] is modified by changing $X^{[\gamma]}$ and $\langle z, n(\gamma)\rangle$ in Part (iv) to $X^{[2n(\gamma)]}$ and $\langle z, 2n(\gamma)\rangle$, respectively, and by adding a Part (v) which states

(v) If $\gamma \subseteq \alpha$, $i = |\gamma|$, and $R_i$ is $R^X_{e,n}$, then $X \neq [e]^{D^{[n]}}$ and $X^{[2n(\gamma)+1]}$ is finite.

The proof of the new Part (v) of the lemma goes as follows. By the inductive hypothesis, it is only necessary to prove (v) for $\gamma = \alpha$. Let $R_{e+1}$ be $R^X_{i,n}$ and let $s_\alpha$ be as in the statement of the lemma. By the argument for Part (iv) of the lemma given in [7], no $Q_\beta$ puts a number into $X^{[2n(\alpha)+1]}$ after stage $s_\alpha$. Suppose that $X = [i]^{D^{[n]}}$. Then, $\lim_s l'(e+1,s) = \infty$. We claim that

$$X^{[2n(\alpha)+1]} = X^{[2n(\alpha)+1]}_{s_\alpha} \cup \{\langle z, 2n(\alpha)+1\rangle | \langle z, 2n(\alpha)+1\rangle \geq R(\alpha) \wedge z \in K\}. \quad (4)$$

To see the right-to-left inclusion of (4), suppose that $x = \langle z, 2n(\alpha)+1\rangle \geq R(\alpha)$ and $z \in K$. Take $s \geq s_\alpha$ such that $s$ is an $\alpha$-stage, $z \in K_s$, $s > x$ and $l'(e+1,s) \geq x$. Since $R(\alpha,[s,e+1]) = R(\alpha)$, $x$ is enumerated into $X$ at stage $[s,e+2]$. For the other inclusion, suppose that $x \in X^{[2n(\alpha)+1]} - X^{[2n(\alpha)+1]}_{s_\alpha}$. Then, $x$ did not enter $X$ through the action of any $Q_\beta$, so $x$ entered $X$ at Step 1.5 of Stage $[s,e+2]$, where $s$ is an $\alpha$-stage $> s_\alpha$. Since $R(\alpha,[s,e+1]) = R(\alpha)$, $x \in \{\langle z, 2n(\alpha)+1\rangle | \langle z, 2n(\alpha)+1\rangle \geq R(\alpha) \wedge z \in K\}$.

From (4), we now get $K \leq_{tt} X^{[2n(\alpha)+1]} \leq_{tt} X \leq_{tt} D^{[n]}$, contradicting assumption that $D^{[n]}$ is Turing-incomplete.

Thus, $X \neq [i]^{D^{[n]}}$, so $l'(e+1,s)$ is bounded as $s \to \infty$. This implies that the set of elements put into $X^{[2n(\alpha)+1]}$ by $R_\alpha$ is finite. As already mentioned, the set of elements put into $X^{[2n(\alpha)+1]}$ by $Q_\beta$ requirements is finite. Thus, $X^{[2n(\alpha)+1]}$ is finite. This completes the proof of Part (v).

There are a few other modifications to the proof of Lemma 3.3 of [7] which are necessary to prove this lemma, but they are routine and are left to the reader. ∎

An ideal $\mathbf{I}$ of $\mathcal{C}_{tt}$ is called $\Sigma^0_3$ if $\{i | deg_{tt}(W_i) \in \mathbf{I}\}$ is $\Sigma^0_3$.

**Theorem 2** *Let* $\mathbf{I}$ *be a* $\Sigma^0_3$*-ideal of* $\mathcal{C}_{tt}$ *consisting of only Turing-incomplete truth-table degrees. Then,* $\mathbf{I}$ *has an exact pair* $\mathbf{b}, \mathbf{c}$ *such that neither of* $\mathbf{b}$ *and* $\mathbf{c}$ *belongs to* $\mathbf{I}$.

**Proof:** This theorem follows from Lemma 1 as Theorem 3.2 follows from Lemma 3.3 in [7]. ∎

**Theorem 3** *If* $\mathbf{a}$ *is a Turing-incomplete c.e. tt-degree, then* $\mathbf{a}$ *is branching.*

**Proof:** Let $\mathbf{a}$ be a Turing-incomplete c.e. truth-table degree and define $\mathbf{I} = \{\mathbf{d} \in \mathcal{C}_{tt} | \mathbf{d} \leq \mathbf{a}\}$. Fix a c.e. set $A \in \mathbf{a}$. We have

$$deg_{tt}(W_i) \in \mathbf{I} \Leftrightarrow (\exists e)(\forall x)(\forall t)(\exists s \geq t)[\{e\}_s(x)\downarrow \wedge W_{i,s}(x) = [e]^{A_s}(x)],$$

so $\mathbf{I}$ is a $\Sigma^0_3$-ideal. All the members of $\mathbf{I}$ are Turing-incomplete. Thus, by Theorem 2, there is an exact pair $\mathbf{b}, \mathbf{c}$ for $\mathbf{I}$ such that neither $\mathbf{b}$ nor $\mathbf{c}$ belongs to $\mathbf{I}$. It follows that $\mathbf{a} < \mathbf{b}, \mathbf{c}$ and that $\mathbf{a}$ is the meet of $\mathbf{b}$ and $\mathbf{c}$ in $\mathcal{C}_{tt}$, so $\mathbf{a}$ is branching. ∎

# 3 Turing Complete Truth-Table Degrees

In this section, we present the main result of the paper, namely, that every Turing-complete tt-incomplete c.e. degree is branching.

**Theorem 4** *If* $\mathbf{b}$ *is a Turing-complete, tt-incomplete c.e. tt-degree, then* $\mathbf{b}$ *is branching in* $\mathcal{C}_{tt}$.

**Proof:** Let $B$ be a c.e. set in $\mathbf{b}$. We want to construct c.e. sets $A_0, A_1$ such that $A_i \not\leq_{tt} B$ for $i = 0, 1$ and such that $deg_{tt}(A_0 \oplus B)$ and $deg_{tt}(A_1 \oplus B)$ have meet $\mathbf{b}$ in $\mathcal{C}_{tt}$. We will actually meet the following requirements $R_{e,i}$ for $e \in \omega$ and $i = 0, 1$:

$$R_{e,i} : [e]^{A_i \oplus B} \neq A_{1-i}.$$

These requirements certainly imply $A_i \not\leq_{tt} B$ for $i = 0, 1$ and the meet requirements will hold automatically due to the way we achieve the $R_{e,i}$ requirements.

An intuitive and oversimplified description of how we meet requirement $R_{e,i}$ is the following. Because $K \not\leq_{tt} B$, we know that if we make $A_i$ look like $\omega$ (above higher priority restraint) and we make $A_{1-i}$ look like $K$ (on a row), then there must be an $x$ such that $[e]^{A_i \oplus B}(x) \neq A_{1-i}(x)$. Not only is there such an $x$, but $K$ can find it; that is, there is a function $f \leq_T K$ such that if $A_i$ looks like $\omega$ up to $f(e)$ (above higher priority restraint) and $A_{1-i}$ looks like $K$ on a specified row up to $f(e)$, then $[e]^{A_i \oplus B} \neq A_{1-i}$. Since $K \leq_T B$, there is a $j_0$ with $\{j_0\}^B = f$. Then, for all $x$, $\{j_0\}^{B_s}_s(x) \to f(x)$ as $s \to \infty$. At stage $s$ of the construction, we will make $A_i$ look like $\omega$ up through $\{j_0\}^{B_s}_s(e)$ and $A_{1-i}$ look like $K_s$ through the same number (on a row). In fact, if we get a new computation $\{j_0\}^{B_s}_s(e)$ at stage $s$, we make $A_i$ look like $\omega$ and $A_{1-i}$ look like $K$ up through $s$. Because of the usual convention that $\{j_0\}^{B_s}_s(e) \leq s$, this suffices to meet $R_{e,i}$. The reason we put potentially many more numbers than necessary to meet $R_{e,i}$ into $A_i$ is to ensure that meet requirements hold.

In outline, the argument that $deg_{tt}(A_0 \oplus B)$ and $deg_{tt}(A_1 \oplus B)$ have meet $deg_{tt}(B) = \mathbf{b}$ in $\mathcal{C}_{tt}$ is as follows. It suffices to show that, given $e_0, e_1$ with $\{e_0\}, \{e_1\}$ total, there is an $e'$ with $\{e'\}$ total such that for all $x$, $[e']^B(x)$ is either

$[e_0]^{A_0 \oplus B}(x)$ or $[e_1]^{A_1 \oplus B}(x)$. Given $x$, $[e']$ finds $M = \max\{|[e_0](x)|, |[e_1](x)|\}$ and then at stage $M$ finds the highest priority $R_{e,i}$ such that $\{j_0\}_M^{B_M}(e)$ converges by an incorrect computation. Then, at some later stage, every number greater than the current restraint on $R_{e,i}$ and less than $M$ will be dumped into $A_i$ and no number smaller than the current restraint will be put in $A_i$. This allows $[e']$ to correctly compute $[e_i]^{A_i \oplus B}(x)$. We will see that this is in fact a truth-table reduction.

To begin the formal construction, we define

$$f(\langle e, m, k\rangle) = \mu y[\{e\}(\langle y, m\rangle)\uparrow \vee [e]^{(\omega^{[\geq m]} \cup F_k) \oplus B}(\langle y, m\rangle)\downarrow \neq K(y)].$$

Since $\omega^{[\geq m]} \cup F_k$ is recursive and $K \not\leq_{tt} B$, $f$ is total. Since $B$ is Turing complete, $f \leq_T B$, say $f = \{j_0\}^B$. The extra parameters in the definition of $f$, compared to the intuitive discussion just given, are to account for the effect of higher priority requirements. If the requirements of higher priority than $R_{e,i}$ control the first $m$ rows and $F_k$ is the set of numbers that the higher priority requirements put into $\omega^{[<m]}$, then $\omega^{[\geq m]} \cup F_k$ is the value $A_i$ would have if $R_{e,i}$ puts every element it can into $A_i$.

We make the following assumptions:

$$\begin{array}{l}\text{if } x = \langle e, m, k\rangle \text{ and } \{e\}(\langle f(x), m\rangle)\downarrow, \text{ then } u(B; j_0, x) > \\ |[e](\langle f(x), m\rangle)| \text{ and } K(f(x)) = K_{u(B;j_0,x)}(f(x));\end{array} \tag{5}$$

$$\{j_0\}_0^D(y)\uparrow \text{ for all } D, y; \tag{6}$$

$$\begin{array}{l}\text{if } \{j_0\}_s^{B_s}(x)\downarrow \text{ and } B_s \restriction u(B_s; j_0, x, s) \neq B_{s+1} \restriction u(B_s; j_0, x, s), \\ \text{then } \{j_0\}_{s+1}^{B_{s+1}}(x)\uparrow.\end{array} \tag{7}$$

The last of these assumptions is the usual "hat trick". With these assumptions, we have the following lemma, which summarizes how we will meet the $R_{e,i}$ requirements.

**Lemma 5** *Let $\{e\}$ be total and let $s$ be such that $\{j_0\}_s^{B_s}(\langle e, m, k\rangle)\downarrow$ and*

$$B_s \restriction u(B_s; j_0, \langle e, m, k\rangle, s) = B \restriction u(B_s; j_0, \langle e, m, k\rangle, s).$$

*If*

$$A_i^{[\geq m]} \restriction s = \omega^{[\geq m]} \restriction s, \tag{8}$$

$$A_i^{[<m]} = F_k, \text{ and} \tag{9}$$

$$(\forall y \leq s)(\langle y, m\rangle \in A_{1-i} \leftrightarrow y \in K_s) \tag{10}$$

*then $R_{e,i}$ is met.*

**Proof:** Let $x = \langle e, m, k\rangle$ and $u = u(B_s; j_0, x, s)$. By the usual convention, $s \geq u, f(x)$ and by the assumption of the lemma, $u = u(B; j_0, x)$. Because $\{e\}$ is total, we have, by definition of $f$, that $[e]^{(\omega^{[\geq m]} \cup F_k) \oplus B}(\langle f(x), m\rangle)\downarrow \neq K(f(x))$. Thus, the lemma will be proven if we can show that $[e]^{A_i \oplus B}(\langle f(x), m\rangle) = [e]^{(\omega^{[\geq m]} \cup F_k) \oplus B}(\langle f(x), m\rangle)$ and that $K(f(x)) = A_{1-i}(\langle f(x), m\rangle)$. Towards establishing the first of these equalities, note that by (5), $s \geq u > |[e](\langle f(x), m\rangle)|$. Thus, (8) and (9) give us the desired equality, since $A_i$ equals $\omega^{[\geq m]} \cup F_k$ up to the length of the appropriate truth table. For the second equality, we have, by (10), $A_{1-i}(\langle f(x), m\rangle) = K_s(f(x))$. Since $s \geq u$, (5) gives $K_s(f(x)) = K(f(x))$, completing the proof. ∎

Let $(R_n)_{n \in \omega}$ be an effective ordering of the $R_{e,i}$'s. The construction will be a finite injury one. From time to time during the construction, an attack may be begun on a requirement $R_n$. The attack may later be canceled. At most one attack on a given requirement will be active at a time. When an attack is made on $R_n$, two parameters $m_n$ and $x_n$ will be defined. If the attack is later canceled, then these values become undefined.

$R_n = R_{e,i}$ *requires attention* at stage $s+1$ if either

$$R_n \text{ is not under attack at the end of stage } s \tag{11}$$

or

$$\text{there is an attack on } R_n \text{ at the end of stage } s,\ \{j_0\}_s^{B_s}(x_{n,s})\downarrow \text{ and } \{j_0\}_{s-1}^{B_{s-1}}(x_{n,s})\uparrow. \tag{12}$$

**Construction:**

**Stage 0:** Do nothing.

**Stage s+1:** Find the highest priority requirement $R_n$ that requires attention. Let $R_n = R_{e,i}$. Proceed according to which case in the definition of requires attention holds.

*(11) holds:* Begin an attack on $R_n$. Set $m_n = s$ and $x_n = \langle e, s, k\rangle$, where $F_k = A_{i,s}^{[<s]}$. Put $\langle y, m_n\rangle$ into $A_i$ for all $y \leq s$. For all $y \in K_s$, put $\langle y, m_n\rangle$ into $A_{1-i}$.

*(12) holds:* Put $\langle y, z\rangle$ into $A_i$ for all $y, z$ with $m_n \leq z \leq s$ and $y \leq s$. For all $y \in K_s$, put $\langle y, m_n\rangle$ into $A_{1-i}$.

In either case, cancel all attacks on $R_{n'}$ with $n' > n$.

**End of Construction**

**Lemma 6** *For each $n$, $R_n$ requires attention only finitely often and an uncanceled attack is made on $R_n$.*

**Proof:** Suppose that the result is true for all $n' < n$. Take $s_0$ such that no $R_{n'}$ with $n' < n$ requires attention at a stage $> s_0$. If no attack on $R_n$ is in progress at the end of stage $s_0$, one will be made at stage $s_0 + 1$, so there is an attack in progress at the end of stage $s_0 + 1$. This attack will never be canceled and $R_n$ never requires attention via (11) after stage $s_0 + 1$. Let $x_n, m_n$ be the values

associated with the final attack and take $s_1 \geq s_0 + 1$ such that $\{j_0\}_{s_1}^{B_{s_1}}(x_n)\downarrow$ and $B_{s_1}\restriction u(B_{s_1}; j_0, x_n, s_1) = B\restriction u(B_{s_1}; j_0, x_n, s_1)$. Then, $R_n$ does not require attention via (12) after stage $s_1 + 1$, so $R_n$ requires attention only finitely often. ∎

**Lemma 7** *Each $R_n$ is met.*

**Proof:** Let $R_n = R_{e,i}$. We may assume that $\{e\}$ is total, since otherwise the result is immediate. Suppose that the final attack on $R_n$ is begun at stage $s_0 + 1$. Then, the final values of $m_n$ and $x_n$ are $s_0$ and $\langle e, s_0, k\rangle = \langle e, m_n, k\rangle$, respectively, where $F_k = A_{i,s_0}^{[<s_0]}$. Let $s + 1 \geq s_0 + 1$ be the last stage at which $R_n$ requires attention. We will use Lemma 5 applied with $m = m_n$ to show that $R_n$ is met. First, we want to show that $\{j_0\}_s^{B_s}(x_n)\downarrow$ and $B_s\restriction u(B_s; j_0, x_n, s) = B\restriction u(B_s; j_0, x_n, s)$. Suppose not. Let $t > s$ be the first stage at which $\{j_0\}_t^{B_t}(x_n)\downarrow$ by a $B$-correct computation. Then, by the "hat trick" (7), $\{j_0\}_{t-1}^{B_{t-1}}(x_n)\uparrow$, so $R_n$ requires attention at stage $t + 1 > s + 1$, contradicting our assumption.

To show (8), note that if $\langle y, z\rangle \in \omega^{[\geq m_n]}\restriction s$, then $m_n \leq z \leq s$ and $y \leq s$. We may have either (11) or (12) holding at stage $s + 1$ (depending on whether $s = s_0$ or $s > s_0$), but in either case, $\langle y, z\rangle$ is put into $A_i$ at stage $s + 1$. Thus, $A_i^{[\geq m_n]}\restriction s = \omega^{[\geq m_n]}\restriction s$.

Towards showing the last two facts we need in order to apply Lemma 5, note that by construction, if some $R_{n'}$ puts a number $y$ into $A_0 \cup A_1$ at stage $s' + 1$, then $y \in \omega^{[\geq m_{n',s'}]} \cap \omega^{[\leq s']}$. Also, each $R_{n'}$ with $n' > n$ is canceled at stage $s_0 + 1$ and any value assigned to $m_{n'}$ at a later stage will be $> s_0$. Furthermore, no $R_{n'}$ with $n' < n$ receives attention at a stage $\geq s_0 + 1$. It follows from these facts that $F_k = A_{i,s_0}^{[<s_0]} = A_i^{[<s_0]} = A_i^{[<m_n]}$, so (9) holds.

Finally, to establish (10), suppose that $y \in K_s$. Then, $\langle y, m_n\rangle \in A_{1-i,s} \subseteq A_{1-i}$ because $R_n$ receives attention at stage $s+1$. Conversely, if $\langle y, m_n\rangle \in A_{1-i}$, then by the considerations of the previous paragraph, $\langle y, m_n\rangle$ was put into $A_{1-i}$ by $R_n$. Since $R_n$ last acts at stage $s + 1$, if $\langle y, m_n\rangle \in A_{1-i}$, then $y \in K_s$. Thus, (10) holds and by Lemma 5, $R_n$ is met. ∎

**Lemma 8** *If $\{e_0\}$ and $\{e_1\}$ are both total, then there is an $e'$ such that $\{e'\}$ is total and for all $x$, either $[e']^B(x) = [e_0]^{A_0 \oplus B}(x)$ or $[e']^B(x) = [e_1]^{A_1 \oplus B}(x)$. (So, if $[e_0]^{A_0 \oplus B} = [e_1]^{A_1 \oplus B} = g$, then $[e']^B = g$.)*

**Proof:** Given $x$, we compute the truth table $[e'](x)$ as follows: Find the truth tables $[e_0](x)$ and $[e_1](x)$ and let $M = \max\{|[e_0](x)|, |[e_1](x)|\}$. Carry out the construction through the end of stage $s_0 = M + 1$. Suppose that at the end of stage $s_0$ there are attacks in progress on requirements $R_0, \ldots, R_{n_0}$. (Note that by construction, if there is an attack in progress on $R_n$ at the end of a stage $s$ and $n' < n$, then there is an attack in progress on $R_{n'}$ at the end of stage $s$.) The truth table $[e'](x)$ will have length $M$ and will query all the numbers

$0, \ldots, M-1$. Given $\sigma \in 2^M$, find the minimum $n \leq n_0$, if any, such that either $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\uparrow$ or $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\downarrow$ and $B_{s_0-1}\restriction u(B_{s_0-1}; j_0, x_{n,s_0}, s_0-1) \neq \sigma\restriction u(B_{s_0-1}; j_0, x_{n,s_0}, s_0-1)$. (Note that if $n \leq n_0$ and $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\downarrow$, then $u(B_{s_0-1}; j_0, x_{n,s_0}, s_0-1) < s_0-1 = M = |\sigma|$.) If no such $n$ exists, then $[e']^\sigma(x)$ is $[e_0]^{A_{0,s_0}\oplus\sigma}(x)$. If such a minimal $n$ exists, then let $R_n$ be $R_{e,i}$. We define $[e']^\sigma(x)$ to be $[e_i]^{(A_{i,s_0}^{[<m_{n,s_0}]}\cup\omega^{[\geq m_{n,s_0}]})\oplus\sigma}(x)$. Then, $e'$ is certainly a truth table reduction. We have to show that $[e']^{B\restriction M}(x)$ is either $[e_0]^{A_0\oplus B}(x)$ or $[e_1]^{A_1\oplus B}(x)$.

Suppose first that there is no $n \leq n_0$ such that either $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\uparrow$ or $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\downarrow$ and, letting $u = u(B_{s_0-1}; j_0, x_{n,s_0}, s_0-1)$, $B_{s_0-1}\restriction u \neq (B\restriction M)\restriction u = B\restriction u$. Then, by induction on $s \geq s_0+1$, it follows that at no such stage $s$ does any $R_n$ with $n \leq n_0$ receive attention. Thus, $A_{0,s_0}^{[<s_0]} = A_{0,s_0}^{[\leq M]} = A_0^{[\leq M]}$. It follows that $A_{0,s_0}\restriction M = A_0\restriction M$ and hence $[e']^{B\restriction M}(x) = [e_0]^{A_{0,s_0}\oplus(B\restriction M)}(x) = [e_0]^{A_0\oplus B}(x)$, as desired.

Now suppose that $n$ is minimal such that $n \leq n_0$ and either $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\uparrow$ or $\{j_0\}_{s_0-1}^{B_{s_0-1}}(x_{n,s_0})\downarrow$ and, letting $u = u(B_{s_0-1}; j_0, x_{n,s_0}, s_0-1)$, $B_{s_0-1}\restriction u \neq (B\restriction M)\restriction u = B\restriction u$. Then, no $R_{n'}$ with $n' < n$ requires attention at any stage $s \geq s_0+1$, so $A_{j,s_0}^{[<m_{n,s_0}]} = A_j^{[<m_{n,s_0}]}$, for $j = 0, 1$. Let $R_n$ be $R_{e,i}$. At some stage $t \geq s_0+1$, $R_n$ will receive attention. At stage $t$, all numbers $\langle y, z\rangle$ with $y < t$ and $m_{n,s_0} \leq z < t$ are put into $A_i$. Since $t \geq s_0+1 > M$, $A_i^{[\geq m_{n,s_0}]}\restriction M = \omega^{[\geq m_{n,s_0}]}\restriction M$. Thus,

$$\begin{aligned} A_i\restriction M &= A_i^{[<m_{n,s_0}]}\restriction M \cup A_i^{[\geq m_{n,s_0}]}\restriction M \\ &= A_{i,s_0}^{[<m_{n,s_0}]}\restriction M \cup \omega^{[\geq m_{n,s_0}]}\restriction M = (A_{i,s_0}^{[<m_{n,s_0}]}\cup\omega^{[\geq m_{n,s_0}]})\restriction M. \end{aligned}$$

Hence, $[e']^{B\restriction M}(x) = [e_i]^{(A_{i,s_0}^{[<m_{n,s_0}]}\cup\omega^{[\geq m_{n,s_0}]})\oplus(B\restriction M)}(x) = [e_i]^{A_i\oplus B}(x)$, as desired. ∎

This completes the proof of Theorem 4.

# 4 Open Problems

We conclude the paper by listing three problems related to the topic of the paper that we are unable to solve using the methods developed here.

Q1) Does every $\Sigma_3^0$-ideal $\mathbf{I}$ of $\mathcal{C}_{tt}$, different from $\mathcal{C}_{tt}$, have an exact pair $\mathbf{b}, \mathbf{c}$ such that neither of $\mathbf{b}$ and $\mathbf{c}$ belongs to $\mathbf{I}$?

(The restriction in (Q1) that $\mathbf{b}, \mathbf{c}$ do not belong to $\mathbf{I}$ is relevant only when $\mathbf{I}$ is a principal ideal. If $\mathbf{I}$ is not principal, then neither half of an exact pair for $\mathbf{I}$ can belong to $\mathbf{I}$.) Theorem 2 gives an affirmative answer to (Q1) when $\mathbf{I}$ does not contain any Turing-complete degrees. Theorem 4 gives an affirmative answer

to (Q1) when $\mathbf{I}$ is a principal ideal and contains Turing complete degrees. An affirmative answer to (Q1) for arbitrary $\mathbf{I}$ would make it possible to obtain our result that every $tt$-incomplete c.e. truth-table degree is branching in $\mathcal{C}_{tt}$ from a more general uniformity of $\mathcal{C}_{tt}$. The method of Section 3 does not appear to give an affirmative answer to (Q1) in the remaining open case, namely, when $\mathbf{I}$ is nonprincipal and contains Turing-complete degrees.

A set $A$ is said to be *bounded truth-table (btt) reducible* to a set $B$ if there are numbers $e$ and $k$ such that $A = [e]^B$ and for all $x$, the truth table $[e](x)$ queries at most $k$ numbers.

Q2) Is every $btt$-incomplete c.e. bounded truth-table degree branching in $\mathcal{C}_{btt}$?

The method of Section 2, unchanged, shows that every Turing-incomplete c.e. bounded truth-table degree is branching in $\mathcal{C}_{btt}$. However, the proof in Section 3 does not show that every Turing-complete btt-incomplete bounded truth-table degree is branching in $\mathcal{C}_{btt}$. This is because in Lemma 8, the truth-table reduction $[e']$ is not bounded, even if the reductions $[e_0]$ and $[e_1]$ are bounded.

So far, when a structural property of $\mathcal{C}_{tt}$ has been shown, it has turned out to be possible to show the same result, by a more difficult proof, for $\mathcal{D}_{tt}(\leq \mathbf{0}'_{tt})$ and then for the structure $\mathcal{D}_{wtt}(\leq \mathbf{0}'_{wtt})$, where $wtt$ stand for weak truth-table reducibility (i.e., Turing reducibility with a recursive bound on the use function). In fact, no elementary difference is known between these three structures. In view of the results in this paper, it is natural to ask

Q3) Is every degree in $\mathcal{D}_{tt}(\leq \mathbf{0}'_{tt})$, other than $\mathbf{0}'_{tt}$, branching in $\mathcal{D}_{tt}(\leq \mathbf{0}'_{tt})$?

A negative answer to this question would be the most interesting, but would be surprising. We believe that the method from Section 3, with additional complications, can be used to give an affirmative answer to (Q3) when the degree is Turing-complete, but the situation for Turing-incomplete degrees is unknown – the exact pair result from Section 2 is not easily adapted to this case.

# References


[1] P. F. Cohen, *Weak truth-table reducibility and the pointwise ordering of 1-1 recursive functions*, Ph.D. thesis, University of Illinois at Urbana-Champaign, 1975.

[2] S. D. Denisov, *On m-degrees of recursively enumerable sets*, Algebra i Logika **9** (1970), 422–427 (Russian), English transl. in Algebra and Logic **9** (1970) 254–256.

[3] Y. L. Ershov and I. A. Lavrov, *The uppersemilattice $L(\gamma)$*, Algebra i Logika **12** (1973), 167–189, 243–244 (Russian), English transl. in Algebra and Logic **12** (1973) 93–106.

[4] Peter A. Fejer, *The density of the nonbranching degrees*, Ann. Pure Appl. Logic **24** (1983), 113–130.

[5] Peter A. Fejer and Richard A. Shore, *A direct construction of a minimal recursively enumerable truth-table degree*, Recursion Theory Week (Proceedings of a Conference Held in Oberwolfach, FRG, March 19–25, 1989) (Berlin) (K. Ambos-Spies, G. H. Müller, and G. E. Sacks, eds.), Lecture Notes in Mathematics, Number 1132, Springer-Verlag, 1990, pp. 187–204.

[6] Alistair H. Lachlan, *Lower bounds for pairs of recursively enumerable degrees*, Proc. London Math. Soc. (3) **16** (1966), 537–569.

[7] André Nies and Richard A. Shore, *Interpreting true arithmetic in the theory of the r.e. truth table degrees*, Ann. Pure Appl. Logic **75** (1995), 269–311.

[8] Piergiorgio Odifreddi, *Classical Recursion Theory, volume 2*, In preparation.

[9] Theodore A. Slaman, *The density of infima in the recursively enumerable degrees*, Ann. Pure Appl. Logic **52** (1991), 155–179.

[10] Robert I. Soare, *Recursively enumerable sets and degrees: The study of computable functions and computably generated sets*, Perspectives in Mathematical Logic, Ω-Series, Springer-Verlag, Berlin, 1987.